arXiv:1605.02490v1 [math.DS] 9 May 2016

ASYMPTOTIC DISTRIBUTION OF VALUES OF ISOTROPIC QUADRATIC FORMS AT S-INTEGRAL POINTS

JIYOUNG HAN, SEONHEE LIM, KEIVAN MALLAHI-KARAI

Abstract. We prove a generalization of a theorem of Eskin-Margulis-Mozes [6] in an S-arithmetic setup: suppose that we are given a finite set of places S over $\mathbb{Q}$ containing the archimedean place, an irrational isotropic form $\mathbf{q}$ of rank $n \geq 4$ on $\mathbb{Q}_S$, a product of p-adic intervals I_p, and a product Ω of star-shaped sets.

We show that if the real part of $\mathbf{q}$ is not of signature $(2,1)$ or $(2,2)$, then the number of S-integral vectors $\mathbf{v} \in \mathsf{T}\Omega$ satisfying simultaneously $\mathbf{q}(\mathbf{v}) \in I_p$ for $p \in S$ is asymptotically $\lambda(\mathbf{q},\Omega)|\mathsf{I}| \cdot \|\mathsf{T}\|^{n-2}$, as T goes to infinity, where $|\mathsf{I}|$ is the product of Haar measures of the p-adic intervals I_p.

Contents

1. Introduction 1
1.1. The Quantitative Oppenheim Conjecture 2
1.2. The S-arithmetic Oppenheim Conjecture 3
1.3. Statements of results 3
2. Preliminaries 7
2.1. Norms on exterior products 7
2.2. Quadratic forms 7
2.3. Orthogonal groups 7
3. The S-arithmetic Geometry of Numbers 8
3.1. S-lattices and a generalization of the α function 8
3.2. Integration on subvarieties 12
4. Passage to homogeneous dynamics 12
5. Counting Results 16
6. Equidistribution of translated orbits 20
7. Establishing upper bounds for the certain integrals 22
7.1. Coarsely superharmonic functions 28
8. Counterexamples for ternary forms 32
Appendix A. A version of Witt theorem 34
References 36

1. Introduction

The Oppenheim conjecture, settled by Margulis in 1986 [14], states that for any non-degenerate indefinite quadratic form $\mathbf{q}$ over $\mathbb{R}^n$, $n \geq 3$, the set $\mathbf{q}(\mathbb{Z}^n)$ of values of integral vectors is a dense subset of $\mathbb{R}$ if $\mathbf{q}$ is *irrational*, i.e. if $\mathbf{q}$ is not proportional to a form with rational coefficients. Both indefiniteness and irrationality conditions are easily seen to be necessary. Margulis' proof

is based on a study of orbits of certain unipotent one-parameter groups on the space of lattices in $\mathbb{R}^3$. The idea of using unipotent flows dates back in an explicit form to Raghunathan and in implicit forms to Cassels and Swinnerton-Dyers. More precisely, Raghunathan observed that Oppenheim conjecture follows from the assertion that every bounded orbit of the orthogonal group $\mathrm{SO}(2,1)$ on $\mathrm{SL}_3(\mathbb{R})/\mathrm{SL}_3(\mathbb{Z})$ is compact. Raghunathan further conjectured in mid-seventies that if G is a connected Lie group, Γ a lattice in G, and U a unipotent subgroup of G, then the closure of any orbit Ux, for $x \in G/\Gamma$ is itself an orbit Lx, where L is a closed connected subgroup of G containing U. Note that the subgroup L depends on x. Raghunathan's conjecture was later proved by Ratner.

1.1. **The Quantitative Oppenheim Conjecture.** Quantitative Oppenheim conjecture, a refinement of Oppenheim conjecture, is about the asymptotic distribution of the values of $\mathbf{q}(\mathbf{v})$, where $\mathbf{v}$ runs over integral vectors in a large ball. To be more precise, let Ω be a radial set defined by $\Omega = \{\mathbf{x} \in \mathbb{R}^n : ||\mathbf{x}|| < \rho(\mathbf{x}/||\mathbf{x}||)\}$, where ρ is a positive continuous function on the unit sphere and $T\Omega$ is the dilation of Ω by a factor $T > 0$. Let us also denote by $\mathbf{N}_{(a,b),\mathbf{q},\Omega}(T)$ the number of vectors $\mathbf{x} \in T\Omega$ such that $\mathbf{q}(\mathbf{x}) \in (a,b)$.

In [3], Dani and Margulis obtained the asymptotic exact lower bound: for $\mathbf{q}$ an irrational indefinite quadratic form on $\mathbb{R}^n$ with $n \geq 3$, for a given interval (a,b) and a set Ω, we have

$$\liminf_{T\to\infty} \frac{\mathbf{N}_{(a,b),\mathbf{q},\Omega}(T)}{\lambda_{\mathbf{q},\Omega}(b-a)T^{n-2}} \geq 1.$$

Note that the denominator is the asymptotic volume of the set $\{\mathbf{v} \in T\Omega : \mathbf{q}(\mathbf{v}) \in (a,b)\}$. The proof of the above theorem is based on studying the distribution of the shifted orbits of the form $u_t K x$, where u_t is a certain one-parameter unipotent subgroup, K is a maximal compact subgroup of the orthogonal group $\mathrm{SO}(\mathbf{q})$ associated to $\mathbf{q}$, and $x \in \mathrm{SL}_n(\mathbb{R})/\mathrm{SL}_n(\mathbb{Z})$. A suitable choice of a function f on $\mathbb{R}^n$ allows one to approximate the number of the integral vectors $\mathbf{v}$ in $T\Omega$ satisfying $\mathbf{q}(\mathbf{v}) \in (a,b)$ by an integral of the form

$$\int_K \widetilde{f}(u_t k x) d\mu(x),$$

where $d\mu$ is the invariant probability measure on $\mathrm{SL}_n(\mathbb{R})/\mathrm{SL}_n(\mathbb{Z})$, and $\widetilde{f} : \mathrm{SL}_n(\mathbb{R})/\mathrm{SL}_n(\mathbb{Z})$ is the Siegel transform of f defined by $\widetilde{f}(g\Gamma) = \sum_{\mathbf{v}\in g\mathbb{Z}^n} f(\mathbf{v})$. Although $\widetilde{f}$ is an unbounded function to which the uniform version of Ratner's theorem proved in [3] does not apply, it is yet possible to establish asymptotically sharp lower bounds by approximating $\widetilde{f}$ from below by bounded functions.

The question of the upper bound, which turned out to be substantially subtler, was taken up in the groundbreaking work [6]: if $\mathbf{q}$ is a form of signature (r,s) with $r \geq 3$ and $s \geq 1$, then

$$\lim_{T\to\infty} \frac{\mathbf{N}_{(a,b),\mathbf{q}}(T)}{\lambda_{\mathbf{q},\Omega}(b-a)T^{n-2}} = 1.$$

For quadratic forms of signature $(2,1)$ and $(2,2)$, certain quadratic forms were constructed in [6] so that by choosing Ω to be the unit ball, for any $\epsilon > 0$,

$$\mathbf{N}_{(a,b),\mathbf{q}_i}(T_j) \gg T_j^i (\log T_j)^{1-\epsilon}$$

for an infinite sequence $T_j \to \infty$. These counterexamples are very well approximable by rational forms. When $\mathbf{q}$ has signature $(2,2)$, it is proven in [7] that if forms of a class called EWAS are excluded, then a similar asymptotic statement continues to hold.

1.2. **The S-arithmetic Oppenheim Conjecture.** The generalization we study in this article involves considering various places. Let S_f be a finite set $\{p_1, \dots, p_s\}$ of odd prime numbers and $S = \{\infty\} \cup S_f$. Each element of S (S_f, respectively) is called a *place* (a finite place, respectively). For $p \in S_f$, the p-adic norm on $\mathbb{Q}$ is defined by $|x|_p = p^{-v_p(x)}$, where $v_p(x)$ is the p-adic valuation of x. The completion of $\mathbb{Q}$ with respect to $|\cdot|_p$ is the field of p-adic numbers and is denoted by $\mathbb{Q}_p$. We will sometimes write $\mathbb{Q}_\infty$ for $\mathbb{R}$. When $p \in S_f$, $\mathbb{Q}_p$ contains a maximal compact subring $\mathbb{Z}_p$ consisting of elements with norm at most 1. We also write $\mathbb{U}_p = \mathbb{Z}_p - p\mathbb{Z}_p$ for the set of multiplicative units in $\mathbb{Z}_p$. When $p = \infty$, we set $\mathbb{U}_p = \{\pm 1\}$. We will use the notation $\mathbb{U}_p^n = \mathbb{Z}_p^n - p(\mathbb{Z}_p^n)$ for $p \in S_f$ and $\mathbb{U}_\infty^n = \mathbb{S}^{n-1}$, the set of vectors of length 1 in $\mathbb{R}^n$. λ_p stands for the Haar measure on $\mathbb{Q}_p$ normalized so that $\lambda_p(\mathbb{Z}_p) = 1$. The Haar measure on $\mathbb{Q}_p^n$ and $\mathbb{Q}_S^n$ are denoted by λ_p^n, λ_S^n, respectively.

We denote by $\mathbb{Q}_S = \prod_{p\in S} \mathbb{Q}_p$ the direct product of $\mathbb{Q}_p$, $p \in S$ and write $\mathbb{Z}_S = \{x \in \mathbb{Q} : |x|_p \le 1,\ \forall p \in S_f\}$ for the ring of S-adic integers. We also denote by λ_S the product measure $\otimes_{p\in S}\lambda_p$.

A quadratic form $\mathbf{q}$ on $\mathbb{Q}_S^n$ is an S-tuple $(\mathbf{q}_p)_{p\in S}$, where $\mathbf{q}_p$ is a quadratic form on $\mathbb{Q}_p$. The form $\mathbf{q}$ is called *isotropic*, if each $\mathbf{q}_p$ is isotropic, that is, if there exists a non-zero vector $\mathbf{v} \in \mathbb{Q}_p^n$ such that $\mathbf{q}_p(\mathbf{v}) = 0$. A quadratic form over $\mathbb{R}$ is isotropic if and only if it is indefinite. $\mathbf{q}$ is called non-degenerate if each $\mathbf{q}_p$ is non-degenerate. Finally, we call $\mathbf{q}$ *rational* if $\mathbf{q} = \lambda\mathbf{q}_0$ for a form $\mathbf{q}_0$ on $\mathbb{Q}_S^n$ and an invertible element $\lambda \in \mathbb{Q}_S$. Otherwise $\mathbf{q}$ is called *irrational*.

Borel and Prasad proved the S-arithmetic Oppenheim conjecture over any number field k. In the case when $k = \mathbb{Q}$, the theorem says that for a non-degenerate irrational isotropic quadratic form $\mathbf{q}$ on $\mathbb{Q}_S^n$ ($n \ge 3$) and any given $\epsilon > 0$, there exists a non-zero vector $\mathbf{x} \in \mathbb{Z}_S^n$ with $|\mathbf{q}_p(\mathbf{x})| < \epsilon$ for all $p \in S$.

1.3. **Statements of results.** The goal of this paper is to prove a quantitative version of the theorem of Borel and Prasad mentioned above. Note that due to the presence of $|S|$ valuations, each one of these objects will depend on $|S|$ parameters.

Let $\mathsf{T} = (T_p)_{p\in S}$, be an S-tuple of positive real numbers with the components $T_p \in p^{\mathbb{Z}}$ for $p \in S_f$. Such an S-tuple T will be called an S-time. The set of all S-time vectors is denoted by $\mathcal{T}_S$. For $\mathsf{T} \in \mathcal{T}_S$, write $\|\mathsf{T}\| = \prod_{p\in S} T_p.$, and $m(\mathsf{T}) = \min_{p\in S} T_p$. For $\mathsf{T} = (T_p), \mathsf{T}' = (T_p') \in \mathcal{T}_S$, define $\mathsf{T} \succeq \mathsf{T}'$ if $T_p \ge T_p'$ for all $p \in S$. Similarly, $\mathsf{T}_i = (T_{p,i})_{p\in S} \to \infty$ means that $T_{p,i} \to \infty$ for each $p \in S$. Since T is of the form $(e^{t_\infty}, p_1^{t_1}, \cdots, p_s^{t_s})$, let us denote $\mathsf{t} = (t_\infty, t_1, \cdots, t_s)$ and consider S-parameter groups

$$\mathbb{T} = \mathbb{R} \times \prod_{p\in S_f} \mathbb{Z}, \qquad \mathbb{T}^+ = \mathbb{R}^+ \times \prod_{p\in S_f} \mathbb{Z}^+.$$

which we will use later. The notions $\mathsf{t} \succeq \mathsf{t}'$ and $\mathsf{t} \to \infty$ are defined accordingly.

For $p \in S$, let $\rho_p : \mathbb{U}_p^n \to \mathbb{R}$ be a positive continuous function. For $p \in S_f$, we will assume throughout the paper that ρ satisfies the following condition:

($\mathbf{I}_\rho$) $$\rho(u\mathbf{x}) = \rho(\mathbf{x}), \quad \forall \mathbf{x} \in \mathbb{U}_p^n, \quad \forall u \in \mathbb{U}_p.$$

Define

(1) $$\Omega = \prod_{p\in S} \Omega_p \subseteq \mathbb{Q}_S^n,$$

where Ω_p is the set of vectors $\mathbf{v}_p$ whose norm is bounded by the value of ρ_p in the direction of $\mathbf{v}_p$. Condition ($\mathbf{I}_\rho$) is indeed very mild and is satisfied by many sets of interest (e.g., the unit ball, which is defined by the constant function). Having fixed an S-time T, we denote by

$\mathsf{T}\Omega = \{(z_p\mathbf{v}_p)_{p\in S} : z_p \in \mathbb{Q}_p, |z_p| \le T_p, \mathbf{v}_p \in \Omega_p\}$ the dilation of Ω by T. For a vector $\mathbf{v} \in \mathbb{Q}_S^n$, we denote $\|\mathbf{v}\| = (\|\mathbf{v}_p\|_p)_{p\in S}$. A p-adic interval of length p^{-b} is a set of the form $I_p = a + p^b\mathbb{Z}_p$, where $a \in \mathbb{Q}_p$ and $b \in \mathbb{Z}$. An S-adic interval I is a product of p-adic interval I_p, with $p \in S$. Let us denote $|\mathsf{I}| = \prod \lambda_p(I_p)$.

Definition 1.1 (Counting and volume functions)**.** *Fixing Ω as above, the counting and volume functions are defined respectively by*

$$\mathbf{N}_{\mathsf{I},\mathbf{q},\Omega}(\mathsf{T}) = \mathrm{card}\{\mathbf{v} \in \mathbb{Z}_S^n \cap \mathsf{T}\Omega : \mathbf{q}_p(\mathbf{v}_p) \in I_p, \forall p \in S\},$$
$$\mathbf{V}_{\mathsf{I},\mathbf{q},\Omega}(\mathsf{T}) = \mathrm{vol}\{\mathbf{v} \in \mathbb{Q}_S^n \cap \mathsf{T}\Omega : \mathbf{q}_p(\mathbf{v}_p) \in I_p, \forall p \in S\},$$

where card *and* vol *denote the cardinality and the Haar measure λ_S^n.*

We start with the following statement about the asymptotic volume of the set of vectors in $T_p\Omega_p \subseteq \mathbb{Q}_p^n$ with the condition $\mathbf{q}(\mathbf{v}) \in \mathsf{I}$. This proposition will be proven in Section 4.

Proposition 1.2. *For a constant $\lambda = \lambda(\mathbf{q}, \Omega)$, we have,*

$$\mathbf{V}_{\mathsf{I},\mathbf{q},\Omega}(\mathsf{T}) \sim \lambda(\mathbf{q}, \Omega) \cdot |\mathsf{I}| \cdot \|\mathsf{T}\|^{n-2},$$

as $\mathsf{T} \to \infty$.

The asymptotic behavior of $\mathbf{N}_{\mathsf{I},\mathbf{q},\Omega}(\mathsf{T})$ is more intricate than the volume asymptotics. The issue for real forms of signature $(2, 1)$ and $(2, 2)$ persists in the S-arithmetic setup. However, p-adic forms of rank 4 behave more nicely than the corresponding real forms of signature $(2, 2)$, which allow us to prove a partial result even in this case.

Definition 1.3. *Let $\mathbf{q} = (\mathbf{q}_p)_{p\in S}$ be an isotropic quadratic form on $\mathbb{Q}_S^n$. We say that $\mathbf{q}$ is exceptional if either $n = 3$, or $n = 4$ and $\mathbf{q}_\infty$ is of signature $(2, 2)$.*

Note that the set $\mathcal{O}_S(n)$ of non-degenerate quadratic forms on $\mathbb{Q}_S^n$ can be identified with a Zariski open subset of $\prod_{p\in S} \mathrm{sym}_n(\mathbb{Q}_p)$, where $\mathrm{sym}_n(\mathbb{Q}_p)$ denotes the set of $n \times n$ matrices over $\mathbb{Q}_p^n$. In this correspondence, a form $\mathbf{q} = (\mathbf{q}_p)_{p\in S}$ is associated to its Gram matrix.

Theorem 1.4. *Let $\mathcal{D}$ be a compact subset of non-exceptional quadratic forms over $\mathbb{Q}_S^n$ of discriminant 1, I an S-interval, and Ω as in (1). Then there exists a constant $C = C(\mathcal{D}, \mathsf{I}, \Omega)$ such that for any $\mathbf{q} \in \mathcal{D}$ and any sufficiently large S-time T we have*

$$\mathbf{N}_{\mathsf{I},\mathbf{q},\Omega}(\mathsf{T}) \le C\|\mathsf{T}\|^{n-2}.$$

Note that this theorem is effective in the sense that the constant C can be explicitly given in terms of the data. The next theorem provides asymptotically sharp bounds for $\mathbf{N}_{\mathsf{I},\mathbf{q},\Omega}(\mathsf{T})$. For the sake of simplicity the theorem below is stated for an individual irrational form. Modifications of the proof, along the lines of the proofs in [6], can be made to establish a uniform version when $\mathbf{q}$ runs over a compact set of forms.

Theorem 1.5. *Let $n \ge 4$ and $\mathbf{q}$ be an irrational isotropic form in $\mathcal{O}(n)$, I be an S-interval, and Ω be as in (1). If $\mathbf{q}$ is non-exceptional, then*

$$\lim_{\mathsf{T}\to\infty} \frac{\mathbf{N}_{\mathsf{I},\mathbf{q},\Omega}(\mathsf{T})}{\lambda(\mathbf{q}, \Omega)|\mathsf{I}| \cdot \|\mathsf{T}\|^{n-2}} = 1.$$

Remark 1.6. From the proof of Theorem 1.5, we can also obtain a partial result in the case when $\mathbf{q}_\infty$ has signature $(2, 2)$. Assume that $\mathsf{T} \to \infty$ while being contained in the cone

$$\log T_\infty \le \delta \sum_{p\in S_f} \log T_p$$

where δ is the constant in Theorem 1.7. Then

$$\frac{\mathbf{N}'_{\mathsf{I},\mathbf{q}}(\mathsf{T})}{\hat{\lambda}(\mathbf{q})|\mathsf{I}|\cdot\|\mathsf{T}\|^{n-2}}\sim 1,$$

where $\mathbf{N}'_{\mathsf{I},\mathbf{q}}(\mathsf{T})$ counts the number of integral solutions of

$$\|\mathbf{v}_\infty\|\le T_\infty,\quad \|\mathbf{v}_p\|_p=T_p,\quad \mathbf{q}(g\mathbf{v})\in I$$

and $\hat{\lambda}(\mathbf{q})$ is a constant depending on $\mathbf{q}$. Note that the conditions above defines the intersection of a ball in the archimedean place with spheres in non-archimedean places.

The proof of Theorem 1.5 rests upon a number of ingredients. We will first relate the counting question to a question about the asymptotic behavior of integrals of the form

$$\int_{\mathsf{K}} f(a_{\mathsf{t}}\mathsf{k}x)dm(\mathsf{k}).$$

Denote the space of unimodular lattices by $\mathcal{L}_S$ (see Definition 3.1 for S-lattices).

Theorem 1.7. *With the notation as above, assume that* $\mathbf{q}$ *is not exceptional. For* $0<s<2$ *and for any* S*-lattice* $\Delta\in\mathcal{L}_S$*,*

$$\sup_{\mathsf{t}>1}\int_{\mathsf{K}}\alpha(a_{\mathsf{t}}\mathsf{k}\Delta)^s dm(\mathsf{k})<\infty.$$

Moreover, the bound is uniform as Δ *varies over a compact subset* $\mathcal{C}$ *of* $\mathcal{L}_S$*. When the signature of* $\mathbf{q}_\infty$ *is* $(2,2)$*, the same inequality is valid if* t *is inside the cone*

$$\mathsf{t}_\infty<\delta\sum_{p\in S_f}\mathsf{t}_p,$$

where δ *is a constant depending on* S *and quadratic form* $\mathbf{q}$*.*

We will deduce the main theorem from the following result:

Theorem 1.8. *Let* $\phi:\mathsf{G}/\Gamma\to\mathbb{R}$ *be a continuous function, and* ν *be a positive continuous function on* $\prod_{p\in S}\mathbb{U}_p^n$*. Assume that for some* $0<s<2$*, we have* $|\phi(\Delta)|<C\alpha(\Delta)^s$ *for all* $\Delta\in\mathsf{G}/\Gamma$*. Let* $x_0\in\mathsf{G}/\Gamma$ *be such that* $\mathsf{H}x_0$ *is not closed. Then*

$$\lim_{\mathsf{t}\to\infty}\int_{\mathsf{K}}\phi(a_{\mathsf{t}}\mathsf{k}x_0)\nu(\mathsf{k})dm(\mathsf{k})=\int_{\mathsf{G}/\Gamma}\phi(y)d\mu(y)\int_{\mathsf{K}}\nu\,dm(\mathsf{k}).$$

The same holds when the signature of $\mathbf{q}_\infty$ *is* $(2,2)$*, assuming that* t *is inside a the cone defined by*

$$\mathsf{t}_\infty<\delta\sum_{p\in S_f}\mathsf{t}_p,$$

where δ *is the same constant as in Theorem 1.7.*

As in [6], the proof of this theorem relies upon a uniform version of Ratner theorem. To explain this version, we will recall some definitions. Let $\mathbf{G}$ be a $\mathbb{Q}$-algebraic group, and let S_f be a set of odd primes. Set $S=S_f\cup\{\infty\}$ as before, and let $\mathsf{G}=\mathbf{G}(\mathbb{Q}_S)=\prod_{p\in S}\mathbf{G}(\mathbb{Q}_p)$. Let Γ be an S-arithmetic subgroup of G, i.e. a group commensurable to $\mathbf{G}(\mathbb{Z}_S)$. Note that $\mathbf{G}(\mathbb{Q}_p)$ is naturally embedded in G. For each $p\in S$, let $U_p=\{u_p(z_p):z_p\in\mathbb{Q}_p\}$ be a one-parameter unipotent $\mathbb{Q}_p$-subgroup, which means that $u_p:\mathbb{Q}_p\to\mathbf{G}(\mathbb{Q}_p)$ is a non-trivial $\mathbb{Q}_p$-rational homomorphism. [17]. We will need a uniform version of Ratner's theorem for the action of unipotent groups on G/Γ, in the special case that Γ is an S-arithmetic subgroup. The following class of groups introduced by Tomanov plays an important role in this theorem.

Definition 1.9. [23] *A connected $\mathbb{Q}$-algebraic subgroup $\mathbf{P}$ of $\mathbf{G}$ is a subgroup* of class $\mathcal{F}$ relative to S *if for each proper normal $\mathbb{Q}$-algebraic subgroup $\mathbf{Q}$ of $\mathbf{P}$, there exists $p \in S$ such that $(\mathbf{P}/\mathbf{Q})(\mathbb{Q}_p)$ contains a non-trivial unipotent element.*

Let Γ be an S-arithmetic lattice in G. If $\mathbf{P}$ is a subgroup of class $\mathcal{F}$ in $\mathbf{G}$, then for any subgroup P' of finite index in $\mathbf{P}(\mathbb{Q}_S)$, $P' \cap \Gamma$ is an S-arithmetic lattice in P'. We will use the following S-arithmetic version of Dani-Margulis theorem:

Theorem 1.10. [13] *Let $\mathbf{G}$ be a $\mathbb{Q}$-algebraic group, $\mathsf{G} = \mathbf{G}(\mathbb{Q}_S)$, and Γ an S-arithmetic lattice in G, and μ denote the G-invariant probability measure on G/Γ. Let $\mathsf{U} = \{(u_p(z_p))_{p\in S} | z_p \in \mathbb{Q}_p\}$ be a one-parameter unipotent $\mathbb{Q}_S$-subgroup of G, and let $\phi : \mathsf{G}/\Gamma \to \mathbb{R}$ be a bounded continuous function. Let $\mathcal{K}$ be a compact subset of G/Γ, and let $\epsilon > 0$. Then there exist finitely many proper subgroups $\mathbf{P}_1, \dots, \mathbf{P}_k$ of $\mathbf{G}$ of class $\mathcal{F}$, and compact subsets $C_i \subseteq X(\mathsf{P}_i, U)$, where $\mathsf{P}_i = \mathbf{P}_i(\mathbb{Q}_S)$, $1 \le i \le k$, such that the following holds: given a compact subset $\mathcal{F}$ of $\mathcal{K} - \cup_i C_i\Gamma/\Gamma$, for all $x \in \mathcal{F}$ and all T with $m(\mathsf{T}) \gg 0$, the following holds*

$$\left| \frac{1}{\lambda_S(\mathsf{I}(\mathsf{T}))} \int_{\mathsf{I}(\mathsf{T})} \phi(u_{\mathsf{z}} x) d\lambda_S(\mathsf{z}) - \int_{\mathsf{G}/\Gamma} \phi d\mu \right| \le \epsilon.$$

It is natural to inquire what happens in the case of exceptional forms. In this direction, we can prove the following theorem:

Theorem 1.11. *Let $I \subseteq \mathbb{Q}_S$ be an interval and Ω be the product of unit balls in $\mathbb{Q}_p^3$, for $p \in S$. Then, there exists an isotropic quadratic form on $\mathbb{Q}_S^3$ which is not proportional to a rational form, a constant $c > 0$, and a sequence $\mathsf{T}_i \to \infty$ such that Then*

$$\mathbf{N}_{I,\mathbf{q}}(\mathsf{T}_j) > c\|\mathsf{T}_j\|(\log \|\mathsf{T}_j\|)^{1-\epsilon}.$$

Since scalar multiples of a single form $\mathbf{q}$ cover all possible equivalence classes of isotropic quadratic forms over $\mathbb{Q}_p^3$ (this can be easily seen from the proof of Proposition 2.1), the above theorem shows that the counterexamples can be constructed in any equivalence class of quadratic forms. This theorem, however, leaves out the case of forms with four variables. It would be interesting to study these forms in details in the future.

Remark 1.12. In this paper, sans serif roman letters are reserved for S-adic objects. Algebraic groups will be denoted by bold letters. For instance, when $\mathbf{G} = \mathrm{SL}_n$, then $\mathbf{G}(\mathbb{Q}_p) = \mathrm{SL}_n(\mathbb{Q}_p)$ and $\mathsf{G} = \prod_{p\in S} \mathbf{G}(\mathbb{Q}_p)$. Element of G, H, etc. are denoted by lower case g, h, etc.

Remark 1.13. Prime powers p^n can be considered both real and p-adic numbers. In some occasions (see Lemma 4.1) we need to transform a real $p^n \in \mathbb{R}$ to $p^n \in Q_p$, which will be denoted by: $(p^n)^\diamond = p^{-n} \in \mathbb{Q}_p$. This notation also renders some formulae more familiar; for instant, the unit p-adic vector in the same direction as a vector $\mathbf{v}$ is $\mathbf{v}/\|\mathbf{v}\|^\diamond$. When $p = \infty$, we set $x^\diamond = x$. For an S-vector $x = (x_p)_{p\in S} \in \mathbb{R}^S$, we set $x^\diamond = (x_p^\diamond)_{p\in S} \in \mathbb{Q}^S$.

Remark 1.14. For simplicity, we have dealt only with the field of rational numbers, but a generalization to the case of number fields is not too difficult.

Acknowledgement. We would like to thank G. Margulis, A. Mohammadi and A. Eskin for helpful discussions. We also acknowledge the support of Seoul National University, MSRI and Jacobs University during our visits. We were informed that Rex Chung has some result on the upper bound but we do not know the details.

2. Preliminaries

In this section, we recall some definitions about $\mathbb{Q}_p$-vector spaces and the quadratic forms defined on them.

2.1. **Norms on exterior products.** Let $1 \leq i \leq n$. One equips the ith exterior product $\bigwedge^i \mathbb{R}^n$ with an inner product defined by

$$\langle \mathbf{x}_1 \wedge \cdots \wedge \mathbf{x}_i, \mathbf{y}_1 \wedge \cdots \wedge \mathbf{y}_i \rangle = \det\left(\mathbf{x}_k \cdot \mathbf{y}_j\right)_{1 \leq k,j \leq i}.$$

For $\mathbf{v} \in \bigwedge^i \mathbb{R}^n$, the induced norm is given by $\|\mathbf{v}\| = |\langle \mathbf{v}, \mathbf{v} \rangle|^{1/2}$. Denote by $\mathbf{e}_1, \ldots, \mathbf{e}_n$ the canonical basis for $\mathbb{R}^n$. For $J = \{1 \leq j_1 < \ldots < j_i \leq n\}$, denote $\mathbf{e}_J = \mathbf{e}_{j_1} \wedge \cdots \wedge \mathbf{e}_{j_i}$. As J runs over all subsets of $\{1, 2, \ldots, n\}$ of i elements, the set $\{\mathbf{e}_J\}$ forms an orthonormal basis for $\bigwedge^i \mathbb{R}^n$ and hence $\|\sum_J a_J \mathbf{e}_J\| = \left(\sum_J a_J^2\right)^{1/2}$. The inner product defined above is preserved by the action of the orthogonal group $\mathrm{O}(n)$ on $\mathbb{R}^n$.

The most convenient norm to work with, for our purpose, is given by $\|\mathbf{v}\| = \max_J |a_J|_p$. The action of $\mathrm{SL}_n(\mathbb{Z}_p)$ on $\bigwedge^i \mathbb{Q}_p^n$ preserves this norm. Indeed, since the entries of $k \in \mathrm{SL}_n(\mathbb{Z}_p)$ are p-adic integers, we immediately have the inequality $\|k\mathbf{v}\|_p \leq \|\mathbf{v}\|_p$. Equality follows by applying this inequality to k^{-1} instead of k.

2.2. **Quadratic forms.** In this subsection, we review the classification of quadratic forms over the p-adic fields $\mathbb{Q}_p$ for odd prime p following [20]. Recall that a quadratic form q on $\mathbb{R}^n$ has signature (r, s) if there is $g \in \mathrm{GL}_n(\mathbb{R})$ such that

$$q(\mathbf{x}) = \mathbf{q}(g\mathbf{x}),$$

where $\mathbf{q}(\mathbf{x}) = \sum_{i=1}^{r} x_i^2 - \sum_{i=r+1}^{r+s} x_j^2$. Two quadratic forms over the field $\mathbb{Q}_p$ are equivalent if and only if they have the same rank, the same discriminant, and the same Hasse invariant. Recall that the discriminant of a non-degenerate quadratic form over a field F is the image of the determinant of the Gram matrix G in the group $F^*/(F^*)^2$. This group contains 4 elements when $F = \mathbb{Q}_p$, and p is an odd prime, representing the parity of $v_p(\det G)$ and whether the image of $\det G$ in the residue field is a quadratic residue or not. Hasse invariant can take two values and is invariant under scalar multiplication. It follows that there are at most 8 different quadratic forms over $\mathbb{Q}_p$ of any given dimension. An isotropic quadratic form $\mathbf{q}$ over $\mathbb{Q}_p$ is called *standard* if it is of the form

$$\mathbf{q}(\mathbf{x}) = x_1 x_n + a_2 x_2^2 + \cdots + a_{n-1} x_{n-1}^2,$$

where $a_i \in \{1, p, u, pu\}$. Here $u \in \mathbb{U}_p$ is a fixed element such that its image in $\mathbb{Z}_p/p\mathbb{Z}_p$ is a non-residue.

2.3. **Orthogonal groups.** For a quadratic form $\mathbf{q} = (\mathbf{q}_p)_{p \in S}$ over $\mathbb{Q}_S$, we define the orthogonal group of $\mathbf{q}$ by $\mathrm{SO}(\mathbf{q}) = \prod_{p \in S} \mathrm{SO}(\mathbf{q}_p)$, where $\mathrm{SO}(\mathbf{q}_p)$ is the p-adic analytic group consisting of matrices of determinant 1 preserving the quadratic form $\mathbf{q}_p$. Let K_∞ be the maximal compact subgroup of $\mathrm{SO}(\mathbf{q}_\infty)$ which is isomorphic to $\mathrm{SO}(r) \times \mathrm{SO}(s)$, where (r, s) is the signature of $\mathbf{q}_\infty$. For $p \in S_f$, we define $K_p = \mathrm{SL}_n(\mathbb{Z}_p) \cap \mathrm{SO}(\mathbf{q}_p)$. It is easy to see that K_p is a maximal subgroup of $\mathrm{SO}(\mathbf{q}_p)$. Finally, we set $\mathsf{K} = \prod_{p \in S} K_p$ and let m_{K}, m_{K_p} be the Haar measures on K, K_p, respectively. Being a subgroups of $\mathrm{SL}_n(\mathbb{Z}_p)$, the linear action of K_p on $\mathbb{Q}_p^n$ preserves the p-adic max norm on $\mathbb{Q}_p^n$. We will also need the following one-parameter subgroups. For each $p \in S_f$,

define

$$a_t^p = \begin{cases} \mathrm{diag}(p^t, 1, \dots, 1, p^{-t}) : & t \in \mathbb{Z} \quad p \in S_f \\ \mathrm{diag}(e^{-t}, 1, \dots, 1, e^t) : & t \in \mathbb{R} \quad p = \infty. \end{cases} \tag{2}$$

We denote by A_p the group consisting of all a_t^p with $t \in \mathbb{Z}$ when $p \in S_f$ and $t \in \mathbb{R}$, when $p = \infty$. Note that these subgroups are isomorphic to $\mathbb{Z}$ when $p \in S_f$ and isomorphic to $\mathbb{R}$, otherwise. Then we can define subgroups of $\mathrm{SO}(\mathbf{q})$ for a standard quadratic form $\mathbf{q}$ as

$$A = \prod A_p = \{a_{\mathtt{t}} = a_{t_\infty}^\infty \cdot \prod_{p \in S_f} a_{t_p}^p\} = \{a_{\mathtt{t}} : \mathtt{t} \in \mathbb{T}\}, \qquad A^+ = \{a_{\mathtt{t}} : \mathtt{t} \in \mathbb{T}^+\}.$$

The following simple fact about ternary forms will be used later.

Proposition 2.1. *Let $\mathbf{q}$ be a non-degenerate isotropic ternary quadratic form over $\mathbb{Q}_p$. Then $\mathrm{SO}(\mathbf{q})$ is isomorphic to $\mathrm{PSL}_2(\mathbb{Q}_p)$.*

Proof. This is well-known for $\mathbb{R}$, so assume $p \in S_f$. The conjugation action of $\mathrm{SL}_2(\mathbb{Q}_p)$ on the space of 2×2 matrices of trace zero over $\mathbb{Q}_p$ preserves the determinant, which can be viewed as a non-degenerate isotropic ternary quadratic form $\mathbf{q}_0$. The kernel is $\pm I$, hence the orthogonal group of $\mathbf{q}_0$ is isomorphic to $\mathrm{PSL}_2(\mathbb{Q}_p)$. For an arbitrary non-degenerate isotropic ternary quadratic form $\mathbf{q}$ and $c \in \mathbb{Q}_p \backslash \{0\}$, the form $c\mathbf{q}$ has the same orthogonal group as $\mathbf{q}$, and for an appropriate value of c has the same discriminant as $\mathbf{q}_0$. □

The proof of the following fact is deferred to the appendix:

Proposition 2.2. *For given $c_1 \in \mathbb{Q}_p$ and $c_2 \in p^{\mathbb{Z}}$, K_p acts transitively on*

$$\{\mathbf{v}_p \in \mathbb{Q}_p^n : \mathbf{q}(\mathbf{v}_p) = c_1 \text{ and } \|\mathbf{v}_p\|_p = c_2\}.$$

3. The S-arithmetic Geometry of Numbers

In this section, we prove some basic results on the S-arithmetic geometry of numbers. Due to lack of reference in S-arithmetic case, we provide proofs.

3.1. S-lattices and a generalization of the α function.

The proof of Oppenheim conjecture is based on the dynamics of unipotent flow on the space of lattices, thus we need the analogous S-arithmetic notion.

Definition 3.1. *Let V be a free $\mathbb{Q}_S$-module of rank n. A $\mathbb{Z}_S$-module Δ is an* S-lattice *in V if there exist $\mathbf{x}_1, \dots, \mathbf{x}_n \in V$ such that*

$$\Delta = \mathbb{Z}_S\mathbf{x}_1 \oplus \cdots \oplus \mathbb{Z}_S\mathbf{x}_n$$

and V is generated by $\mathbf{x}_1, \dots, \mathbf{x}_n$ as a $\mathbb{Q}_S$-module.

Note that an S-lattice is a discrete and cocompact subgroup of $\mathbb{Q}_S^n$. The base lattice Δ_0 is the S-lattice generated by the canonical basis $\mathbf{e}_1, \dots, \mathbf{e}_n$. Note that $\mathrm{SL}_n(\mathbb{Q}_S)$ acts on the set of lattices in $\mathbb{Q}_S^n$ and the orbit of Δ_0 under this action can be identified with $\mathcal{L}_S := \mathrm{SL}_n(\mathbb{Q}_S)/\mathrm{SL}_n(\mathbb{Z}_S)$. For an S-lattice Δ, we say that a subspace $L \subseteq \mathbb{Q}_S^n$ is Δ-*rational* if the intersection $L \cap \Delta$ is an S-lattice in L, that is, L is a subspace of $\mathbb{Q}_S^n$ generated by finite number of elements in Δ. Suppose that L is i-dimensional and $L \cap \Delta = \mathbb{Z}_S\mathbf{v}^1 \oplus \cdots \oplus \mathbb{Z}_S\mathbf{v}^i$. Define $d_\Delta(L)$ or simply $d(L)$ by

$$d(L) = d_\Delta(L) = \prod_{p \in S} \|\mathbf{v}^1 \wedge \cdots \wedge \mathbf{v}^i\|_p. \tag{3}$$

The definition of $d(L)$ is independent of the choice of basis. This is an analogue of the volume of the quotient space $L/(L \cap \Delta)$ for lattices in $\mathbb{R}^n$. If $L = \{0\}$, we write $d(L) = 1$. We now show that there is indeed a more intimate connection between the $\mathcal{L}_S$ and the space $\mathcal{L}$ of lattices (of covolume 1) in $\mathbb{R}^n$. More precisely, we show that the $\mathcal{L}_S$ fibers over $\mathcal{L}$ in such a way that many natural functions on $\mathcal{L}_S$ factor through the projection map.

Definition 3.2 (The projection map)**.** *Let $\Delta \in \mathcal{L}_S$. Define $\pi(\Delta)$ to be the set of vectors $\mathbf{v} \in \mathbb{R}^n$ for which there exists $\mathbf{w} \in \Delta$ with $\mathbf{w}_\infty = \mathbf{v}$ and $\mathbf{w}_p \in \mathbb{Z}_p^n$ for all $p \in S_f$.*

Remark 3.3. For any vector $\mathbf{v} \in \pi(\Delta)$ there is exactly one vector $\mathbf{w} \in \Delta$ satisfying $\mathbf{w}_\infty = \mathbf{v}$. Indeed, if both $\mathbf{w}_1$ and $\mathbf{w}_2$ satisfy this property, then $\mathbf{w} = \mathbf{w}_1 - \mathbf{w}_2 \in \Delta$ is a non-zero vector with real component zero. This implies that $(\prod_{p \in S_f} p^m)\mathbf{w} \to 0$ as $m \to \infty$, which contradicts the discreteness of Δ.

Proposition 3.4. *$\pi(\Delta)$ is a lattice of covolume 1 in $\mathbb{R}^n$.*

Proof. Since the p-adic components of the elements of $\pi(\Delta)$ belong to compact sets $\mathrm{SL}_n(\mathbb{Z}_p)$, for $p \in S_f$, the vectors $\mathbf{w} = \mathbf{v}_\infty$ cannot have an accumulation point, since then by passing to a subsequence, we can find an accumulation point for a sequence of vectors in Δ.

Start with a $\mathbb{Z}_S$-basis $\mathbf{v}^1, \ldots, \mathbf{v}^n$ for Δ. By replacing the vectors $\mathbf{v}^i$ with certain $\mathbb{Z}_S$ linear combinations of them, we may assume that the determinant of the matrix A_p with columns $\mathbf{v}_p^1, \ldots, \mathbf{v}_p^n$ is a unit in $\mathbb{Z}_p$, for every $p \in S_f$. Note that this is equivalent to the condition

$$\|\mathbf{v}_p^1 \wedge \cdots \wedge \mathbf{v}_p^n\|_p = 1 \tag{4}$$

for every $p \in S_f$. We now claim that $\mathbf{v}_\infty^1, \ldots, \mathbf{v}_\infty^n$ form a $\mathbb{Z}$-basis for $\pi(\Delta)$. If $\mathbf{w} \in \pi(\Delta)$, then $\mathbf{w} = \mathbf{v}_\infty$ for some $\mathbf{v} \in \Delta$ with $\mathbf{v}_p \in \mathbb{Z}_p^n$. Write $\mathbf{v} = \sum_{k=1}^n c_k \mathbf{v}^k$, for $c_k \in \mathbb{Z}_S$, and let $\mathbf{c}$ be the column vector with entries $c_1, \ldots, c_n$. Since $A_p\mathbf{c} = \mathbf{v}_p$, $A_p^{-1} \in \mathrm{SL}_n(\mathbb{Z}_p)$, and $\mathbf{v}_p \in \mathbb{Z}_p^n$, we deduce that $\mathbf{c} \in \mathbb{Z}_p^n$ for all $p \in S_f$. This implies that $\mathbf{c} \in \mathbb{Z}^n$, which proves our claim. Since Δ is unimodular, (4) implies that $\|\mathbf{v}_\infty^1 \wedge \cdots \wedge \mathbf{v}_\infty^n\|_\infty = 1$, showing that $\pi(\Delta)$ has covolume 1. □

Definition 3.5. *For an S-lattice Δ, define*

$$\alpha_i^S(\Delta) = \sup\left\{\frac{1}{d(L)} : L \text{ is a } \Delta\text{-rational subspace of dimension } i\right\},$$

where $1 \le i \le n$ and

$$\alpha^S(\Delta) = \max\left\{\alpha_i(\Delta) : 0 \le i \le n\right\}.$$

The function α^S is not only an analogue of the original function α, but is in fact intimately related to it.

Proposition 3.6. *For any $\Delta \in \mathcal{L}_S$, we have*

$$\alpha^S(\Delta) = \alpha(\pi(\Delta)).$$

Proof. Assume that $\alpha_i^S(\Delta) = 1/d(L)$ where L is the $\mathbb{Z}_S$-span of $\mathbf{v}^1, \ldots, \mathbf{v}^i$. Similar to the proof of Proposition 3.4, after possibly replacing $\mathbf{v}^1, \ldots, \mathbf{v}^i$ by another basis, we can assume that $\|\mathbf{v}_p^1 \wedge \cdots \wedge \mathbf{v}_p^i\|_p = 1$ for all $p \in S_f$. This implies that $d(L) = \|\mathbf{v}_\infty^1 \wedge \cdots \wedge \mathbf{v}_\infty^i\|_\infty$. As before, $\mathbf{v}_\infty^1, \ldots, \mathbf{v}_\infty^i$ form a $\mathbb{Z}$-basis for $\pi(L)$, implying that $\alpha_i(\pi(\Delta)) \ge \alpha_i^S(\Delta)$. The converse can be proven in a similar way. □

Since S is fixed throughout the discussion, we will henceforth drop the superscript S in α_i^S and α^S. We now define an S-adic Siegel transform. Let $f : \mathbb{Q}_S^n \to \mathbb{R}$ be a bounded function vanishing outside a compact set. The S-adic Siegel transform of f is defined as follows: for $g \in \mathrm{SL}_n(\mathbb{Q}_S)$, let $\tilde{f}(g) = \sum_{\mathbf{v}\in\mathbb{Z}_S^n} f(g\mathbf{v})$. More generally, $\tilde{f}$ is defined on the space of S-lattices; for such a lattice Δ, define

$$\tilde{f}(\Delta) = \sum_{\mathbf{v}\in\Delta} f(\mathbf{v}).$$

Since the diagonal embedding of $\mathbb{Z}_S^n$ in $\mathbb{Q}_S$ is discrete, the defining sum involves only finitely many non-zero terms. The following lemma is a natural extension of Schmidt's theorem in geometry of numbers.

Lemma 3.7. *(S-adic Schmidt lemma) Let $f : \mathbb{Q}_S^n \to \mathbb{R}$ be a bounded function vanishing outside a compact subset. Then there exists a positive constant $c = c(f)$ such that $\tilde{f}(\Delta) < c\alpha^S(\Delta)$ for any S-lattice Δ in $\mathbb{Q}_S^n$.*

Proof. Without loss of generality, we can assume that f is the characteristic function of $B = \prod_{p\in S} B_p$, where B_p is a ball of radius p^{n_p}, for $p \in S$ and a ball of radius R for $p = \infty$. Set $D = \prod_{p\in S_f} p^{n_p}$ and consider the scalar map $\delta : \Delta \to \Delta$ defined by $\delta(\mathbf{x}) = D \cdot \mathbf{x}$. Note that if $\mathbf{x} \in \Delta \cap B$, then $\|\mathbf{x}\|_p \le 1$ for all $p \in S_f$ and $\|\mathbf{x}\|_\infty \le RD$. In particular, $\pi(\delta(\mathbf{x}))$ will be inside a ball B' of radius RD. Note that Schmidt's lemma states that

$$\mathrm{card}(\Delta' \cap B') = O(\alpha(\Delta')),$$

holds for any lattice Δ' in $\mathbb{R}^n$, where the implies constant depends on B'. Using this fact and Remark 3.3 we obtain

$$\mathrm{card}(\Delta \cap B) \le \mathrm{card}(\pi(\Delta) \cap B') = O(\alpha(\pi(\Delta)) = O(\alpha^S(\Delta)).$$

□

Lemma 3.8. *Let Δ be an S-lattice in $\mathbb{Q}_S^n$ and let L and M be two Δ-rational subspaces. Then we have that*

$$d(L)d(M) \ge d(L \cap M)d(L + M), \tag{5}$$

where $d = d_\Delta$ is defined in (3).

Proof. We first verify that if L and M are Δ-rational, then $L \cap M$ and $L + M$ are also Δ-rational. Since a subspace of $\mathbb{Q}_S^n$ is Δ-rational if and only if it is generated by elements of Δ, $L + M$ is clearly a Δ-rational subspace of $\mathbb{Q}_S^n$. On the other hand, consider a projection map $\pi : \mathbb{Q}_S^n \to \mathbb{Q}_S^n/\Delta$. Note that the projection image $\pi(H)$ of a subspace H in $\mathbb{Q}_S^n$ is closed if and only if $H \cap \Delta$ is a lattice subgroup in H, that is, H is Δ-rational. Since π is proper, if we set $H = \pi^{-1}(\overline{\pi(L \cap M)}) = \pi^{-1}(\overline{\pi(L) \cap \pi(M)})$, we can easily check that H is Δ-rational and $H = L \cap M$.

Let $p : \mathbb{Q}_S^n \to \mathbb{Q}_S^n/(L\cap M)$. Since $d_\Delta(H) = d_{p(\Delta)}(p(H))d_\Delta(L\cap M)$ for any Δ-rational subspace H, the inequality (5) is equivalent to

$$d_{p(\Delta)}(L)d_{p(\Delta)}(M) \ge d_{p(\Delta)}(L + M).$$

Let $\{\mathbf{v}_1, \dots, \mathbf{v}_\ell\}$ and $\{\mathbf{w}_1, \dots, \mathbf{w}_m\}$ be bases of $p(L)$ and $p(M)$ consisting of elements in $p(\Delta)$, respectively. Then since $(p(L) \cap p(\Delta)) + (p(M) \cap p(\Delta)) \subseteq p(L + M) \cap p(\Delta)$,

$$\begin{aligned} d_{p(\Delta)}(L)d_{p(\Delta)}(M) &= \|\mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_\ell\|\|\mathbf{w}_1 \wedge \cdots \wedge \mathbf{w}_m\| \\ &\ge \|\mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_\ell \wedge \mathbf{w}_1 \wedge \cdots \wedge \mathbf{w}_m\| \ge d_{p(\Delta)}(L + M). \end{aligned}$$

□

The following Lemma is the S-arithmetic version of Lemma 3.10 in [6].

Lemma 3.9. *The function α belongs to $L^r(\mathrm{SL}_n(\mathbb{Q}_S)/\mathrm{SL}_n(\mathbb{Z}_S))$ for all values of $1 \leq r < n$.*

Proof. Consider the map $\pi : \mathcal{L}_S \to \mathcal{L}$. Denote the invariant probability measures on $\mathcal{L}_S$ and $\mathcal{L}$ by μ_S and μ, respectively. We claim that π is measure-preserving, that is, for any measurable $A \subseteq \mathcal{L}_{\mathbb{R}}$, we have $\mu_S(\pi^{-1}(A)) = \mu(A)$. To see this, define $\nu(A) = \mu_S(\pi^{-1}(A))$. One can easily see that ν is an $\mathrm{SL}_n(\mathbb{R})$-invariant probability measure on $\mathcal{L}$, implying that $\nu = \mu$. The result will immediately follows from Proposition 3.6. □

We can now prove the following generalization of the classical Siegel's formula in geometry of numbers.

Proposition 3.10. *(S-arithmetic Siegel integral formula) Let f be a continuous function with compact support on $\mathbb{Q}_S^n$, $\mathsf{G} = \mathrm{SL}_n(\mathbb{Q}_S)$ and $\Gamma = \mathrm{SL}_n(\mathbb{Z}_S)$. Then*

$$\int_{\mathsf{G}/\Gamma} \widetilde{f} d\mu = \int_{\mathbb{Q}_S^n} f d\lambda_S^n + f(0).$$

The proof is based on the following lemma. For each $p \in S$, we denote by λ_p^n the Haar measure on $\mathbb{Q}_p^n$ normalized such that $\lambda_p^n(\mathbb{Z}_p^n) = 1$. Denote by δ_p the delta measure at the zero point of $\mathbb{Q}_p^n$.

Lemma 3.11. *Let ν be an $\mathrm{SL}_n(\mathbb{Q}_S)$-invariant Radon measure on $\mathbb{Q}_S^n$. Then ν is a linear combination of the measures of the form $\bigotimes_{p\in S} \nu_p$, where each ν_p is either the Haar measure λ_p^n or the delta measure δ_p.*

Proof. Observe that if ν is a positive Radon measure on a locally compact space X invariant under the action of a group H and aH has a finite number of orbits, namely $X_1, \ldots, X_m$, then ν naturally decomposes into the sum $\nu = \sum_{j=1}^m \nu_i$, where $\nu_i(A) = \nu(A \cap X_i)$ is H-invariant and supported on X_i, for $1 \leq i \leq m$. Let us now consider the case at hand, namely, the action of $\mathrm{SL}_n(\mathbb{Q}_S)$ on $\mathbb{Q}_S^n$. Since the natural action of $\mathsf{G} = \mathrm{SL}_n(\mathbb{Q}_p)$ on $\mathbb{Q}_p^n$ has two orbits on $\mathbb{Q}_p^n$, the G-orbits in the action on $\mathbb{Q}_S^n$ are of the form $\prod_{p\in S} X_p$, where X_p is either the zero point or $\mathbb{Q}_p^n - \{0\}$. Each orbit is a homogenous space G/L, where L is the stabilizer of an arbitrary point in the respective orbit. The invariant measure on the homogeneous space G/L, if exists, is unique up to a factor. The result follows. □

Proof of Proposition 3.10. Note that by Lemmas 3.7 and 3.9, $\widetilde{f}$ is integrable. This implies that the map $f \mapsto \int_{\mathsf{G}/\Gamma} \widetilde{f} d\mu$ defines a positive linear functional on the space of compactly supported continuous functions on $\mathbb{Q}_S^n$, hence defines a Radon measure ν on $\mathbb{Q}_S^n$. Since μ is G-invariant, so is ν. Using Lemma 3.11, ν splits as a linear combination of measures of the form $\bigotimes_{p\in S} \nu_p$, where each ν_p is either λ_p^n or the delta measure δ_p. The coefficient of $\bigotimes_{p\in S} \lambda_p^n$ in this linear combination is one. Indeed, taking an increasing sequence of compactly supported non-negative continuous functions f_n approximating the normalized characteristic functions of increasingly larger balls in $\mathbb{Q}_S^n$, one obtains $\widetilde{f_n} \to 1$, and $\int_{\mathbb{Q}_S^n} f_n d\lambda_S^n \to 1$. Similarly, by choosing a family f_n of non-negative continuous functions with support in a ball of radius $1/n$ with $|f_n| \leq 1$ and $f_n(0) = 1$, and applying the Lebesgue's dominant convergence theorem, we have

$$\lim_{n\to\infty} \int_{\mathsf{G}/\Gamma} \widetilde{f_n} d\mu = 1,$$

which implies that the coefficient of $f(0)$ is 1. It remains to be shown that the rest of the coefficients are zero. This can be obtained in a similar way from the fact that a lattice Δ in $\mathbb{Q}_S^n$ cannot contain a non-zero point $(\mathbf{v}_p)_{p\in S}$ with $\mathbf{v}_{p_0} = 0$ for some $p_0 \in S$. Otherwise, one can find an appropriate sequence n_i of S-integers, such that $|n_i|_q \to 0$ for all $q \neq p_0$ and hence $n_i\mathbf{v} \to 0$ contradicting the discreteness of Δ. From here, and by choosing appropriate sequences of continuous functions as above, we can deduce that the rest of the coefficients are zero. □

3.2. **Integration on subvarieties.** Now let us introduce volume forms of submanifolds in $\mathbb{Q}_S^n$ inherited from the volume form $d\lambda_S^n$ of $\mathbb{Q}_S^n$ in order to measure the volumes of orbits of maximal compact subgroups of orthogonal groups in $\mathbb{Q}_S^n$. Since such maximal compact subgroups are products of maximal compact subgroups at each place, it suffices to deal with volume forms of p-adic submanifolds in $\mathbb{Q}_p^n$.

Let us introduce two equivalent definitions of volume form on submanifolds. Recall that any orbit of the p-adic maximal compact subgroup $K_p = \mathrm{SL}_n(\mathbb{Z}_p) \cap \mathrm{SO}(\mathbf{q}_p)$ of $\mathrm{SO}(\mathbf{q}_p)$ for a given quadratic form $\mathbf{q}_p$ on $\mathbb{Q}_p^n$ is of the form

$$\{\mathbf{w} \in \mathbb{Q}_p^n : ||\mathbf{w}||_p = p^{c_1}, \mathbf{q}_p(\mathbf{w}) = c_2\},$$

where $c_1 \in \mathbb{Z}$ and $c_2 \in \mathbb{Q}_p$ are some constants. Note that the above K_p-orbit is an open subset of the $(n-1)$-dimensional p-adic variety in $\mathbb{Q}_p^n$ given by

$$\{\mathbf{w} \in \mathbb{Q}_p^n : \mathbf{q}_p(\mathbf{w}) = c_2\}.$$

Hence from now on, we will consider Y as a compact open subset of a d-dimensional p-adic variety in $\mathbb{Q}_p^n$ or a parallelepiped $\mathbb{Z}_p\mathbf{v}_1 \oplus \cdots \oplus \mathbb{Z}_p\mathbf{v}_d$ where $\mathbf{v}_1, \ldots, \mathbf{v}_d \in \mathbb{Q}_p^n$. Then Y is contained in $p^{-r}\mathbb{Z}_p^n$ for some $r \in \mathbb{N}$.

Consider the projection $\pi_\ell : p^{-r}\mathbb{Z}_p^n \to p^{-r}\mathbb{Z}_p^n/p^\ell\mathbb{Z}_p^n$ and let $Y_\ell = \pi_\ell(Y)$. Then the volume form ν_d defined over Y is defined by

$$\nu_d(Y) = \lim_{\ell\to\infty} \frac{\#Y_\ell}{p^{d\ell}}.$$

As in the real case, we can define the volume form of a p-adic submanifold by describing the volume form of its tangent space. Let Y be a parallelepiped $\mathbb{Z}_p\mathbf{v}_1 \oplus \cdots \oplus \mathbb{Z}_p\mathbf{v}_d$, where $\mathbf{v}_1, \ldots, \mathbf{v}_d$ are linearly independent in $\mathbb{Q}_p^n$. Then the volume form ν_d' of the parallelepiped is defined as

$$\nu_d'(\mathbb{Z}_p\mathbf{v}_1 \oplus \cdots \oplus \mathbb{Z}_p\mathbf{v}_d) = ||\mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_d||_p,$$

where $||\cdot||$ is the maximum norm of $\bigwedge^d(\mathbb{Q}_p^n)$.

We will use the fact (see [22, Theorem 9]) that the normalized measures ν_d and ν_d' are equal.

4. Passage to homogeneous dynamics

In this section, we define a p-adic analogue of J_f function introduced in Section 3 in [6] for the real case. This function is used to reduce the counting problem to the asymptotic formula of certain integrals on the space of S-lattices. Fix a prime $p \in S_f$, and let $\mathbf{q}$ be a standard quadratic form on $\mathbb{Q}_p^n$. Let $\pi_1 : \mathbb{Q}_p^n \to \mathbb{Q}_p$ be the projection to the first coordinate and $\mathbb{Q}_{p,+}^n$ be the set $\{\mathbf{x} \in \mathbb{Q}_p^n : \pi_1(\mathbf{x}) \neq 0\}$. We denote by $p^{\mathbb{Z}}$ the subset of $\mathbb{Q}_p$ consisting of p-powers.

Lemma 4.1. *Let f be a continuous function of compact support on $\mathbb{Q}_{p,+}^n$ which satisfies the invariance property*

($\mathbf{I}_f$) $$f(ux_1, x_2, \ldots, x_{n-1}, u^{-1}x_n) = f(x_1, x_2, \ldots, x_n)$$

for all units $u \in \mathbb{U}_p$. Let ν be a non-negative continuous function on the unit sphere $\mathbb{U}_p^n$ such that

($\mathbf{I}_\nu$) $$\nu(u\mathbf{x}) = \nu(\mathbf{x}),$$

for all $u \in \mathbb{U}_p$. Let J_f be the real-valued function on $p^{\mathbb{Z}} \times \mathbb{Q}_p \subset \mathbb{Q}_p^2$ defined by

$$J_f(p^{-r}, \zeta) = \frac{1}{p^{r(n-2)}} \int_{\mathbb{Q}_p^{n-2}} f(p^{-r}, x_2, \dots, x_{n-1}, x_n)\ dx_2 \cdots dx_{n-1}, \tag{6}$$

where in the integral, $x_n = p^r\left(\zeta - \mathbf{q}(0, x_2, \dots, x_{n-1}, 0)\right)$. Then for any $\epsilon > 0$, if t and $\|\mathbf{v}\|$ are sufficiently large, we have

$$\left| c(K_p) p^{t(n-2)} \int_{K_p} f(a_t k\mathbf{v}) \nu(k^{-1}\mathbf{e}_1) dm(k) - J_f\left(p^t \|\mathbf{v}\|^\diamond, \zeta\right) \nu\left(\frac{\mathbf{v}}{\|\mathbf{v}\|^\diamond}\right) \right| < \epsilon, \tag{7}$$

where $c(K_p) = \mathrm{vol}(K_p.\mathbf{e}_1)/(1 - 1/p)$ and m is the normalized Haar measure on K_p.

Proof. Let π_i denote the projection onto the ith coordinate in $\mathbb{Q}_p^n$, and $a_t := a_t^p = \mathrm{diag}\left(p^t, 1, \dots, 1, p^{-t}\right)$ as before. Since f has a compact support, for $t \gg 1$, if $f(a_t\mathbf{w}) \neq 0$, we have

$$\|\mathbf{w}\|_p = |\pi_1(\mathbf{w})|_p.$$

This implies that on the support of the function $k \mapsto f(a_t k\mathbf{v})$, for t large enough we have

$$a_t k\mathbf{v} = \left(\|\mathbf{v}\|_p^\diamond p^t \cdot u(k\mathbf{v}), \pi_2(k\mathbf{v}), \dots, \pi_{n-1}(k\mathbf{v}), \mathbf{v}_n u(k\mathbf{v})^{-1}\right). \tag{8}$$

Using ($\mathbf{I}_f$) we have

$$f(a_t k\mathbf{v}) = f\left(\|\mathbf{v}\|_p^\diamond\, p^t, \pi_2(k\mathbf{v}), \dots, \pi_{n-1}(k\mathbf{v}), v_n\right).$$

where v_n is determined by $\mathbf{q}\left(\|\mathbf{v}\|_p^\diamond\, p^t, \pi_2(k\mathbf{v}), \dots, \pi_{n-1}(k\mathbf{v}), v_n\right) = \mathbf{q}(\mathbf{v})$. Since K_p preserves the norm, given $\delta > 0$, for $t > t_0(\delta)$, if $f(a_t k\mathbf{v}) \neq 0$, we have

$$\left| k\mathbf{v}/\|\mathbf{v}\|_p^\diamond - u(\pi_1(k\mathbf{v}))\mathbf{e}_1 \right|_p < \delta$$

for a unit $u(\pi_1(k\mathbf{v})) \in \mathbb{U}_p$. This implies that,

$$\left| \frac{\mathbf{v}}{\|\mathbf{v}\|_p^\diamond} - k^{-1}(u(\pi_1(k\mathbf{v}))\mathbf{e}_1) \right|_p = \left| \frac{\mathbf{v}}{\|\mathbf{v}\|_p^\diamond} - u(\pi_1(k\mathbf{v}))(k^{-1}\mathbf{e}_1) \right|_p < \delta.$$

Since ν is continuous and satisfies ($\mathbf{I}_\nu$), by choosing $\delta > 0$ small enough we can assure that if $t > t_0(\delta)$, and $f(a_t k\mathbf{v}) \neq 0$, then the following holds:

$$\left| \nu(k^{-1}\mathbf{e}_1) - \nu\left(\frac{\mathbf{v}}{\|\mathbf{v}\|_p^\diamond}\right) \right| < \epsilon. \tag{9}$$

For sufficiently large t, if $f(a_t kv) \neq 0$, then the vector $\left(\|\mathbf{v}\|_p^\diamond\, p^t, \pi_2(k\mathbf{v}), \dots, \pi_{n-1}(k\mathbf{v}), v_n\right)$ will be in a bounded distance from $\mathbf{e}_1$.

As the K_p-invariant measure on the orbit $K_p.\mathbf{v}$ is the push-forward of the normalized Haar measure on K, one can approximate the integral in (6) by the integration on the K_p-orbit $K_p.(\|\mathbf{v}\|_p^\diamond\, \mathbf{e}_1) \subset \mathbb{Q}_p^n$. For $t > \max\{t', t''\}$,

$$\left| \int_{K_p} f(a_t k\mathbf{v}) \nu(k^{-1}\mathbf{e}_1) dm(k) \right. \tag{10}$$
$$\left. - \int_{K_p.(\|\mathbf{v}\|_p^\diamond\, \mathbf{e}_1)} f\left(\|\mathbf{v}\|_p^\diamond\, p^t, \pi_2(k\mathbf{v}), \dots, \pi_{n-1}(k\mathbf{v}), v_n\right) \nu\left(\frac{\mathbf{v}}{\|\mathbf{v}\|^\diamond}\right) d\lambda_{n-1} \right|_p \leq \epsilon,$$

where λ_{n-1} is the normalized Haar measure on $K.(\|\mathbf{v}\|_p^\diamond\, \mathbf{e}_1)$. □

Remark that ($\mathbf{I}_\nu$) is an essential condition but ($\mathbf{I}_f$) is not a real restriction since we can easily modify the give function to satisfy ($\mathbf{I}_f$).

Proposition 4.2. *Suppose h is a real-valued continuous function of compact support defined on $(\mathbb{Q}_p^n - \{0\}) \times \mathbb{Q}_p$. Then*

$$\lim_{t\to\infty} \frac{1}{p^{(n-2)t}} \int_{\mathbb{Q}_p^n} h(p^t\mathbf{v}, \mathbf{q}(\mathbf{v}))d\mathbf{v} = \mathrm{vol}(K\cdot\mathbf{e}_1) \sum_{z\in\mathbb{Z}} p^{(n-2)z} \int_K \int_{\mathbb{Q}_p} h(p^{-z}k\mathbf{e}_1, \zeta)d\zeta dm(k).$$

where $\mathrm{vol} = \mathrm{vol}_{n-1}$ *denote the induced* $(n-1)$*-dimensional volume from* $\mathbb{Q}_p^n$.

Proof. Let us make the substitution $\mathbf{w} = p^t\mathbf{v}$. This gives $d\mathbf{w} = 1/p^{nt}d\mathbf{v}$, hence we have

$$\frac{1}{p^{(n-2)t}} \int_{\mathbb{Q}_p^n} h(p^t\mathbf{v}, \mathbf{q}(\mathbf{v}))d\mathbf{v} = \frac{1}{p^{(n-2)t}} \int_{\mathbb{Q}_p^n} h(\mathbf{w}, p^{-2t}\mathbf{q}(\mathbf{w}))p^{nt}d\mathbf{w} = p^{2t} \int_{\mathbb{Q}_p^n} h(\mathbf{w}, p^{-2t}\mathbf{q}(\mathbf{w}))d\mathbf{w}.$$

Replacing p^{-2t} by p^{-t}, the left integral in (11) can be changed to

$$L_1(h) := \lim_{t\to\infty} p^t \int_{\mathbb{Q}_p^n} h(\mathbf{w}, p^{-t}\mathbf{q}(\mathbf{w}))d\mathbf{w}.$$

With the decomposition $\mathbb{Q}_p^n = \bigcup_{z\in\mathbb{Z}} p^{-z}\mathbb{U}_p^n$, write

$$h = \sum_{z\in\mathbb{Z}} h_z, \tag{11}$$

where $h_z(\mathbf{v}, x) = h(\mathbf{v}, x)\mathbf{1}_{p^{-z}\mathbb{U}_p^n}(\mathbf{v})$. Since h has compact support, all but finitely many of h_z are zero. We first claim that for h with $\mathrm{supp}\, h \subseteq \mathbb{U}_p^n \times \mathbb{Q}_p$,

$$\lim_{t\to\infty} \frac{1}{p^{(n-2)t}} \int_{\mathbb{Q}_p^n} h(p^t\mathbf{v}, \mathbf{q}(\mathbf{v}))d\mathbf{v} = \mathrm{vol}(K.\mathbf{e}_1) \int_K \int_{\mathbb{Q}_p} h(k\mathbf{e}_1, \zeta)d\zeta dm(k).$$

Then, for h with $\mathrm{supp}\, h \subseteq p^{-z_0}\mathbb{U}_p^n$ where $z_0 \in \mathbb{Z}$, define $h'(\mathbf{v}, x) = h(p^{-z_0}\mathbf{v}, p^{-2z_0}x)$, which has a compact support in $\mathbb{U}_p^n \times \mathbb{Q}_p$. Using change of variables, we have

$$\begin{aligned}
L_1(h) &= \lim_{t\to\infty} p^t \int_{\mathbb{Q}_p^n} h(\mathbf{v}, p^{-t}\mathbf{q}(\mathbf{v}))d\mathbf{v} = \lim_{t\to\infty} p^t \int_{p^{-z_0}\mathbb{U}_p^n} h(\mathbf{v}, p^{-t}\mathbf{q}(\mathbf{v}))d\mathbf{v} \\
&= p^{nz_0} \lim_{t\to\infty} p^t \int_{\mathbb{U}_p^n} h(p^{-z_0}\mathbf{v}, p^{-t-2z_0}\mathbf{q}(\mathbf{v}))d\mathbf{v} \\
&= p^{nz_0} \lim_{t\to\infty} p^t \int_{\mathbb{U}_p^n} h'(\mathbf{v}, \mathbf{q}(\mathbf{v}))d\mathbf{v} = \mathrm{vol}(K.\mathbf{e}_1) \int_K \int_{\mathbb{Q}_p} h'(k\mathbf{e}_1, \zeta)d\zeta dm(k) \\
&= \mathrm{vol}(K.\mathbf{e}_1)p^{nz_0} \int_K \int_{\mathbb{Q}_p} h(p^{-z_0}k\mathbf{e}_1, p^{-2z_0}\zeta)d\zeta dm(k) \\
&= \mathrm{vol}(K.\mathbf{e}_1)p^{(n-2)z_0} \int_K \int_{\mathbb{Q}_p} h(p^{-z_0}k\mathbf{e}_1, \zeta')d\zeta' dm(k),
\end{aligned}$$

where the last line relies on the change of variables $\zeta' = p^{-2z_0}\zeta$. Now, using the decomposition (11), Proposition 4.2 follows for an arbitrary h.

Now let us prove the claim. Let h satisfy $\operatorname{supp} h \subseteq \mathbb{U}_p^n \times \mathbb{Q}_p$. Furthermore, since we put $\mathbf{q}(\mathbf{v})$ in the second variable and $\mathbf{q}$ is the quadratic form with p-adic integral coefficients, we can add the assumption that $\operatorname{supp} h \subseteq \mathbb{U}_p^n \times \mathbb{Z}_p$.

Let us denote by $\mathcal{A}_k$ the subspace of the space of continuous functions on $\mathbb{U}_p^n \times \mathbb{Z}_p$ which is spanned by the functions of the form $h(\mathbf{v}, x) = f(\mathbf{v})g(x)$, with $f(\mathbf{v}) = \mathbf{1}_A(\mathbf{v})$ and $g(x) = \mathbf{1}_B(x)$, where $A \subseteq \mathbb{U}_p^n$ and $B \subseteq \mathbb{Z}_p$ are respectively the preimages of subsets $\bar{A} \subseteq (\mathbb{Z}/p^k\mathbb{Z})^n$ and $\bar{B} \subseteq \mathbb{Z}/p^k\mathbb{Z}$ by the reduction map red_k. We will also set $\mathcal{A} = \bigcup_{k=1}^{\infty} \mathcal{A}_k$. Using Stone-Weierstrass theorem for p-adic fields ([5]), $\mathcal{A}$ is dense in the space of all continuous functions on $\mathbb{U}_p^n \times \mathbb{Z}_p$. Let us denote the reduction maps $\mathbb{Z}_p \to \mathbb{Z}_p/p^k\mathbb{Z}_p$ (and also $\mathbb{Z}_p^n \to (\mathbb{Z}_p/p^k\mathbb{Z}_p)^n$) by red_k. Let h be a real-valued continuous function of compact support contained in $\mathbb{U}_p^n \times \mathbb{Z}_p$. We will do some reduction steps: we may assume that $h(\mathbf{v}, x) = f(\mathbf{v})g(x)$, and $f(\mathbf{v}) = \mathbf{1}_A(\mathbf{v})$ and $g(x) = \mathbf{1}_B(x)$, with

$$\begin{aligned} A &= \mathrm{red}_k^{-1}(\overline{A}) \subseteq \mathbb{U}_p^n, \\ B &= \mathrm{red}_k^{-1}(\overline{B}) \subseteq \mathbb{Z}_p. \end{aligned} \tag{12}$$

In other words,

$$\mathbf{v} \in A, \mathbf{w} \in \mathbb{Z}_p^n \text{ with } \|\mathbf{w}\| < p^{-k} \implies \mathbf{v} + \mathbf{w} \in A,$$

$$x \in B, y \in \mathbb{Z}_p \text{ with } |y| < p^{-k} \implies x + y \in B.$$

It thus suffices to show that

$$\begin{aligned} L_1(h) &= \lim_{t\to\infty} p^t \mu_n(\{\mathbf{v} \in A, \mathbf{q}(\mathbf{v}) \in p^t B) \\ &= \mathrm{vol}(K.\mathbf{e}_1) m_K(\{k \in K : k\mathbf{e}_1 \in A\}) \mu_1(B). \end{aligned}$$

Let us define

$$\begin{aligned} C_t(A, B) &= \{\mathbf{v} \in \mathbb{Z}_p^n : \mathbf{v} \in A, p^{-t}\mathbf{q}(\mathbf{v}) \in B\}, \\ C_t'(A, B) &= \{\mathbf{u} + p^t\mathbf{w} : \mathbf{u} \in A, \mathbf{q}(\mathbf{u}) = 0, \mathbf{w} \in \mathbb{Z}_p^n, 2\beta(\mathbf{u}, \mathbf{w}) \in B\}. \end{aligned}$$

Here β is the associated bilinear form to the quadratic form $\mathbf{q}$. Recall the polarization identity:

$$\mathbf{q}(\mathbf{v} + \mathbf{w}) = \mathbf{q}(\mathbf{v}) + 2\beta(\mathbf{v}, \mathbf{w}) + \mathbf{q}(\mathbf{w}).$$

We claim that $C_t(A, B) = C_t'(A, B)$ for any $t > k$. First, we show that $C_t'(A, B) \subseteq C_t(A, B)$. Assume that $\mathbf{v} = \mathbf{u} + p^t\mathbf{w}$ with $\mathbf{u} \in A, \mathbf{w} \in \mathbb{Z}_p^n$ satisfying $\mathbf{q}(\mathbf{u}) = 0, 2\beta(\mathbf{u}, \mathbf{w}) \in B$. This implies that

$$\mathbf{q}(\mathbf{u} + p^t\mathbf{w}) = 2p^t\beta(\mathbf{u}, \mathbf{w}) + p^{2t}\mathbf{q}(\mathbf{w})^2 = p^t(2\beta(\mathbf{u}, \mathbf{w}) + p^t\mathbf{q}(\mathbf{w})^2)$$

Since $\mathbf{u} \in A$ and $\|p^t\mathbf{v}\| \leq p^{-t}$, for $t > k$, we have $\mathbf{v} \in A$. Moreover,

$$p^{-t}\mathbf{q}(\mathbf{v}) = 2\beta(\mathbf{u}, \mathbf{w}) + p^t\mathbf{q}(\mathbf{w})^2 \in B.$$

Conversely, if $\mathbf{v} \in C_t(A, B)$, set $\mathbf{v}' = \mathbf{e}_1 + \frac{1}{2}\mathbf{q}(\mathbf{v})\mathbf{e}_n$. Note that $\mathbf{q}(\mathbf{v}') = \mathbf{q}(\mathbf{v})$ and $\|\mathbf{v}\| = \|\mathbf{v}'\| = 1$. Hence by Proposition 2.2, there exists $k \in K$ such that $\mathbf{v} = k\mathbf{v}'$. Set $\mathbf{u} = k\mathbf{e}_1$ and $\mathbf{w} = \frac{1}{2}p^{-t}\mathbf{q}(\mathbf{v})k\mathbf{e}_n$, so that $\mathbf{v}$ decomposes as $\mathbf{v} = k(\mathbf{e}_1 + \frac{1}{2}\mathbf{q}(\mathbf{v})\mathbf{e}_n) = \mathbf{u} + p^t\mathbf{w}$. Furthermore, since $\mathbf{v} \in A$ and $\|\mathbf{u} - \mathbf{v}\| = \|\frac{1}{2}\mathbf{q}(\mathbf{v})k\mathbf{e}_n\| \leq p^{-t} < p^{-k}$, we have $\mathbf{u} \in A$. Clearly, we have $\mathbf{q}(\mathbf{v}) = \mathbf{q}(k\mathbf{e}_1) = 0$ and

$$2\beta(\mathbf{u}, \mathbf{w}) = 2\beta(k\mathbf{e}_1, \frac{1}{2}p^{-t}\mathbf{q}(\mathbf{v})k\mathbf{e}_n) = p^{-t}\mathbf{q}(\mathbf{v})\beta(k\mathbf{e}_1, k\mathbf{e}_n) = p^{-t}\mathbf{q}(\mathbf{v}) \in B.$$

This equality implies that

$$\mu_n(C_t(A,B)) = \mu_n(C_t'(A,B)) = \lim_{\ell\to\infty} \frac{|\mathrm{red}_\ell(C_t'(A,B))|}{p^{n\ell}}.$$

Set $A_\ell = \mathrm{red}_\ell(A) \subseteq \mathbb{Z}/p^\ell\mathbb{Z}$ and $B_\ell = \mathrm{red}_\ell(B) \subseteq \mathbb{Z}/p^\ell\mathbb{Z}$. We will denote the induced quadratic form over $\mathbb{Z}_p/p^\ell\mathbb{Z}_p$ by $\mathbf{q}$ as well. First observe that for $\ell/2 > t > k$, we have

$$\mathrm{red}_\ell(C_t'(A,B)) = \{\overline{u} + p^t\overline{w} : \bar{u} \in A_\ell, \mathbf{q}(\bar{u}) = 0, 2\beta(\bar{u},\bar{w}) \in B_\ell\}.$$

For any choice of $\overline{u} \in A_\ell \cap \{\overline{u} : \mathbf{q}(\overline{u}) = 0\}$ there exist exactly $|B_\ell|$ choices for $\overline{w}$ such that $2\beta(\bar{u},\bar{w}) \in B_\ell$. Hence, in order to find $\mu_n(C_t(A,B))$, it suffices to find the cardinality of the fibers of the map $\Phi : (\mathbb{Z}/p^\ell\mathbb{Z})^n \times (\mathbb{Z}/p^\ell\mathbb{Z})^n \to (\mathbb{Z}/p^\ell\mathbb{Z})^n$ defined by

$$\Phi(\overline{u},\overline{w}) = \overline{u} + p^t\overline{w}.$$

We will show that all the fibers have the same size. If $\Phi(\overline{u}_1,\overline{w}_1) = \Phi(\overline{u}_2,\overline{w}_2)$, then $\overline{u_1} \equiv \overline{u_2} \pmod{p^t}$. Conversely, suppose that $\overline{u_1} \in A_\ell$, $\mathbf{q}(\overline{u_1})$ and $\overline{v} = \overline{u_1} + p^t\overline{w_1}$ for some $\overline{w_1}$ which satisfies $2\beta(\bar{u_1},\bar{w_1}) \in B_\ell$. Let $\overline{u_2} \in A_\ell$ be such that $\mathbf{q}(\overline{u_2}) = 0$, $u_1 \equiv u_2 \pmod{p^t}$. Write $\overline{u_2} - \overline{u_1} = p^t z$. Set $\overline{w_2} = \overline{w_1} - z$, hence $\overline{v} = \overline{u_2} + p^t\overline{w_2}$. First note that $\mathbf{q}(\overline{u_1}) = \mathbf{q}(\overline{u_2}) = 0$ combined with

$$\mathbf{q}(\overline{u_2}) = \mathbf{q}(u_1 + p^t z) = q(u_1) + 2p^t\beta(\overline{u_1}, z) + p^{2t}\mathbf{q}(z)$$

implies that $\beta(\overline{u_1}, z)$ is divisible by p^t. This implies that

$$2\beta(\overline{u_2},\overline{w_2}) = 2\beta(\overline{u_1} - p^t z, \overline{w_1} + z) = 2\beta(\overline{u_1},\overline{w_1}) - 2p^t\beta(z,\overline{w_1}+z) + 2\beta(\overline{u_1}, z) \in B_\ell$$

This shows that the fibers of Φ are in one-to-one correspondence with the equivalence classes of $\overline{u} \in A_\ell$ module p^t. This implies that for all sets A, B satisfying (12), for $\ell \geq t > k$, we have

$$|\mathrm{red}_\ell(C_t(A,B))| = N_t|A_\ell \cap \{\overline{u} : \mathbf{q}(\overline{u}) = 0\}| \times |B_\ell|.$$

Note that N_t does not depend on ℓ. In the special case $A = \mathbb{U}_p^n$ and $B = \mathbb{Z}_p$, $C_t(A,B)$ is precisely the p^{-t}-neighborhood of the set $\{\mathbf{u} \in \mathbb{U}_p^n : \mathbf{q}(\mathbf{u}) = 0\}$. It follows that

$$\lim_{t\to\infty} p^t\mu_n(C_t(\mathbb{U}_p^n,\mathbb{Z}_p)) = \mu_{n-1}(\{\mathbf{u} \in \mathbb{U}_p^n : \mathbf{q}(\mathbf{u}) = 0\}) = \mathrm{vol}(K.\mathbf{e}_1).$$

From here, for general A and B we obtain

$$\begin{aligned}\lim_{t\to\infty} p^t\mu_n(C_t(A,B)) &= \lim_{t\to\infty} p^t\mu_n(C_t(\mathbb{U}_p^n,\mathbb{Z}_p))\frac{\mu_n(C_t(A,B))}{\mu_n(C_t(\mathbb{U}_p^n,\mathbb{Z}_p))} \\ &= \mathrm{vol}(K.\mathbf{e}_1)\lim_{\ell\to\infty}\frac{|A_\ell \cap \{\mathbf{v} \in (\mathbb{Z}/p^\ell\mathbb{Z})^n : \mathbf{q}(\mathbf{v}) = 0\} \cdot |B_\ell|}{|\{\mathbf{v} \in (\mathbb{Z}/p^\ell\mathbb{Z})^n : \mathbf{q}(\mathbf{v}) = 0\} \cdot |\mathbb{Z}/p^\ell\mathbb{Z}|} \\ &= \mathrm{vol}(K.\mathbf{e}_1)m_K\{k \in K : k\mathbf{e}_1 \in A\}\mu_1(B).\end{aligned}$$

□

5. Counting Results

In this section, we prove Theorems 1.4 and 1.5. The proofs depend on an ergodic result which will be proven in the next section. In the course of these proofs we will deal with compactly supported continuous function $h : (\mathbb{Q}_S^n - \{0\}) \times \mathbb{Q}_S \to \mathbb{R}$ of the form $h(\mathbf{v},\zeta) = \prod_{p\in S} h_p(\mathbf{v}_p,\zeta_p)$. For h in this space, define

$$L(h) = \lim_{\mathsf{t}\to\infty} \|\mathsf{T}\|^{-(n-2)} \int_{\mathbb{Q}_S^n} h\left(\frac{\mathbf{v}}{\mathsf{T}^\diamond}, \mathbf{q}(\mathbf{v})\right)\, d\mathbf{v}.$$

Proof of Theorem 1.4. Without loss of generality, we may assume that there exists a compact set $C \subset \mathsf{G}$ so that every quadratic form in $\mathcal{D}$ is of the form $\mathbf{q}^g$ for some $g \in C$, where $\mathbf{q}$ is a fixed standard form, since there are finitely many equivalence classes of quadratic forms over $\mathbb{Q}_S$. Since C is compact, there exists β such that for every $g = (g_p)_{p\in S} \in C$ and every $\mathbf{v} \in \mathbb{Q}_S^n$, $\beta^{-1}\|\mathbf{v}_p\|_p \le \|g_p\mathbf{v}_p\|_p \le \beta\|\mathbf{v}_p\|_p$ for every $p \in S$. Let $\epsilon > 0$ be given and $g \in C$ be arbitrary. Let $f = \prod_{p\in S} f_p$ be a continuous non-negative function of compact support on $\mathbb{Q}_S^n$ such that $J_{f_\infty} \ge (1+\epsilon)^{1/|S|}$ on $[\beta^{-1}, 2\beta] \times I_\infty$ and $J_{f_p} \ge (1+\epsilon)^{1/|S|}$ on $p^{\mathbb{Z}} \cap \{x \in \mathbb{Q}_p : \beta^{-1} \le |x|_p \le 2\beta\} \times I_p$ for $p \in S_f$. It is clear that if $\mathbf{v} \in \mathbb{Q}_S^n$ satisfies the conditions

$$T_\infty \le \|\mathbf{v}_\infty\| \le 2T_\infty, \|\mathbf{v}_p\|_p = T_p, \quad \mathbf{q}(g\mathbf{v}) \in I \tag{13}$$

then $J_f(\|g\mathbf{v}\|^\diamond/|\mathsf{T}|^\diamond, \mathbf{q}(g\mathbf{v})) \ge 1+\epsilon$. By Lemma 4.1 and Lemma 3.6 in [6], for sufficiently large t,

$$c(\mathsf{K})\|\mathsf{T}\|^{n-2}\int_{\mathsf{K}} f(a_\mathsf{t}\mathsf{k}g\mathbf{v})dm(\mathsf{k}) \ge 1$$

if $\mathbf{v}$ satisfies (13). Summing over $\mathbf{v} \in \mathbb{Z}_S^n$, it follows that the number of vectors $\mathbf{v} \in \mathbb{Z}_S^n$ satisfying the conditions (13) is bounded from above by

$$\sum_{\mathbf{v}\in\mathbb{Z}_S} c(\mathsf{K})\|\mathsf{T}\|^{n-2}\int_{\mathsf{K}} f(a_t\mathsf{k}g\mathbf{v})dm(\mathsf{k}) = c(\mathsf{K})\|\mathsf{T}\|^{n-2}\int_{\mathsf{K}} \tilde{f}(a_\mathsf{t}\mathsf{k}g)dm(\mathsf{k}).$$

The statement follows from a geometric sum argument. □

Proposition 5.1. *Let $f_p : \mathbb{Q}_p^n \to \mathbb{R}$ be a positive continuous function with a compact support in $\mathbb{Q}_{p,+}^n$, and $f = \prod_{p\in S} f_p$. Assume further that f_p satisfies the property ($\mathbf{I}_f$) when $p \in S_f$. Let $\nu_p : \mathbb{U}_p^n \to \mathbb{R}$ be positive continuous functions satisfying the condition ($\mathbf{I}_\nu$) and $\nu = \prod_{p\in S}\nu_p$. For a non-exceptional quadratic form $\mathbf{q}$, $\epsilon > 0$ and every $\mathsf{g} \in \mathsf{G}$, there exists a positive S-time t_0 so that for $\mathsf{t} \succ \mathsf{t}_0$,*

$$\left|\|\mathsf{T}\|^{-(n-2)}\sum_{\mathbf{v}\in\mathbb{Z}_S^n} J_f\left(\frac{\|\mathsf{g}\mathbf{v}\|^\diamond}{\mathsf{T}^\diamond}, \mathbf{q}(\mathsf{g}\mathbf{v})\right)\nu\left(\frac{\mathsf{g}\mathbf{v}}{\|\mathsf{g}\mathbf{v}\|^\diamond}\right) - c(\mathsf{K})\int_{\mathsf{K}}\tilde{f}(a_\mathsf{t}\mathsf{k}\mathsf{g})\nu(\mathsf{k}^{-1}\mathbf{e}_1)dm(\mathsf{k})\right| \le \epsilon, \tag{14}$$

where $c(\mathsf{K}) = \prod_{p\in S} c(K_p)$.

Proof. Since $J_f = \prod_{p\in S} J_{f_p}$ has compact support, by Theorem 1.4 for given $\mathsf{g} \in \mathsf{G}$, the number of $\mathbf{v} \in \mathbb{Z}_S^n$ such that $J_f(\mathsf{g}, \mathbf{v}, \mathsf{t})\nu(\mathsf{g}, \mathbf{v}) \ne 0$ is bounded by $c\|\mathsf{T}\|^{n-2}$ for some $c > 0$. Take t_0 such that Lemma 4.1 and Lemma 3.6 in [6] hold for $\mathsf{g}\mathbf{v}$ and ϵ/c. Then for $\mathsf{t} \succ \mathsf{t}_0$,

$$\begin{aligned}
&\left|\|\mathsf{T}\|^{-(n-2)}\sum_{\mathbf{v}\in\mathbb{Z}_S^n} J_f\left(\frac{\|\mathsf{g}\mathbf{v}\|^\diamond}{\mathsf{T}^\diamond}, \mathbf{q}(\mathsf{g}\mathbf{v})\right)\nu\left(\frac{\mathsf{g}\mathbf{v}}{\|\mathsf{g}\mathbf{v}\|^\diamond}\right) - c(\mathsf{K})\int_{\mathsf{K}}\tilde{f}(a_\mathsf{t}\mathsf{k}\mathsf{g})\nu(\mathsf{k}^{-1}\mathbf{e}_1)dm(\mathsf{k})\right| \\
&\le \|\mathsf{T}\|^{-(n-2)}\sum_{\mathbf{v}\in\mathbb{Z}_S^n}\left|J_f\left(\frac{\|\mathsf{g}\mathbf{v}\|^\diamond}{\mathsf{T}^\diamond}, \mathbf{q}(\mathsf{g}\mathbf{v})\right)\nu\left(\frac{\mathsf{g}\mathbf{v}}{\|\mathsf{g}\mathbf{v}\|^\diamond}\right) - c(\mathsf{K})\|\mathsf{T}\|^{n-2}\int_{\mathsf{K}} f(a_\mathsf{t}\mathsf{k}\mathsf{g}\mathbf{v})\nu(\mathsf{k}^{-1}\mathbf{e}_1)dm(\mathsf{k})\right| \\
&\le C(f)e^{-(n-2)t_0}\prod_{p\in S_f} p^{-(n-2)t_p}\cdot\epsilon/c\cdot c\|\mathsf{T}\|^{n-2} = C(f)\epsilon.
\end{aligned}$$

Here $C(f)$ is a constant depending on f which comes from the fact that if $|a_i - b_i| < \epsilon$ and $|a_i| < A, |b_i| < B$, then $|\prod a_i - \prod b_i| < C(A,B)\epsilon$. Note that both terms in the second line of the equations are bounded by a constant depending on f. □

Lemma 5.2. *Let f and ν be as in Theorem 5.1. Let $h_\infty(\mathbf{v}_\infty, \zeta_\infty) = J_{f_\infty}(\|\mathbf{v}_\infty\|_\infty, \zeta_\infty)\nu_\infty(\mathbf{v}_\infty/\|\mathbf{v}_\infty\|_\infty)$, $h_p(\mathbf{v}_p, \zeta_p) = J_{f_p}(\|\mathbf{v}_p\|_p^\diamond, \zeta_p)\nu_p(\mathbf{v}_p/\|\mathbf{v}_p\|_p^\diamond)$ for $p \in S_f$. Set $h(\mathbf{v}, \zeta) = \prod_{p\in S} h_p(\mathbf{v}_p, \zeta_p)$. Then we have*

$$L(h) = c(\mathsf{K}) \int_{\mathsf{G}/\Gamma} \tilde{f}(g)\, dg \prod_{p\in S} \int_{K_p} \nu_p(k_p^{-1}\mathbf{e}_1) dm(k_p)$$

Proof. Since for any unit u,

$$\int_{\mathbb{Q}_p^{n-1}} f(p^{-r}, x_2, \dots, x_{n-1}, x_n) dx_2 \dots dx_n = \int_{\mathbb{Q}_p^{n-1}} f(p^{-r}u, x_2, \dots, x_{n-1}, x_n) dx_2 \dots dx_n,$$

we have that

$$\begin{aligned}\int_{\mathbb{Q}_p^n} f dx_1 \dots dx_n &= \int_{\mathbb{Q}_p} \int_{\mathbb{Q}_p^{n-1}} f(x_1, x_2, \dots x_{n-1}, x_n) dx_2 \dots dx_n \cdot dx_1 \\ &= p^r(1-p^{-1}) \sum_{r\in\mathbb{Z}} \int_{\mathbb{Q}_p^{n-1}} f(p^{-r}, x_2, \dots x_{n-1}, x_n) dx_2 \dots dx_n.\end{aligned}$$

The change of variables $p^{-r}x_n + \mathbf{q}(0, x_2, \dots, x_{n-1}, 0) = \zeta$ yields $p^r dx_n = d\zeta$, and hence

$$\begin{aligned}\int_{\mathbb{Q}_p^n} f dx_1 \dots dx_n &= \sum_{r\in\mathbb{Z}} \int_{\mathbb{Q}_p} \int_{\mathbb{Q}_p^{n-2}} f(p^{-r}, x_2, \cdots, x_{n-1}, x_n) dx_2 \dots dx_{n-1}(1-p^{-1}) d\zeta \\ &= \sum_{r\in\mathbb{Z}} \int_{\mathbb{Q}_p} J_f(p^{-r}, \zeta) p^{r(n-2)}(1-1/p) d\zeta.\end{aligned}$$

The result will now follow from Proposition 3.10), Proposition 4.2, and Lemma 3.9 in [6]. The computations proceeds verbatim those in Lemma 3.9 in [6].

□

Proof of Lemma 1.2. Since $\Omega = \prod_{p\in S} \Omega_p \subset \mathbb{Q}_S^n$, it suffices to show that for any choice of intervals $I_p \subseteq \mathbb{Q}_p$, $p \in S$, there exists a constant $\lambda(\mathbf{q}_p, \Omega_p)$ such that $\mathrm{vol}\{\mathbf{v} \in T_p\Omega_p : \mathbf{q}_p(\mathbf{v}) \in I\}$ is asymptotically $\lambda(\mathbf{q}_p, \Omega_p)T_p^{n-2}$. The case $p = \infty$ is in Lemma 3.8 (ii) of [6]. Hence we may assume that $p \in S_f$, and write $I_p = a + p^b\mathbb{Z}_p$, for $a \in \mathbb{Q}_p$ and $b \in \mathbb{Z}$.

Recall that $\Omega_p = \{\mathbf{v} \in \mathbb{Q}_p^n : \|\mathbf{v}\|_p \le \rho_p(\mathbf{v}/\|\mathbf{v}\|_p^\diamond)\}$, where $\rho(u\mathbf{v}) = \rho(\mathbf{v})$ for any $\mathbf{v} \in \mathbb{U}_p^n$ and $u \in \mathbb{U}_p$. Substituting $g^{-1}\Omega_p$ by Ω_p, we may assume that $\mathbf{q}_p$ is standard. Moreover, the function ρ' determining $g^{-1}\Omega_p$ satisfies $\rho'(\mathbf{v}) = \rho'(u\mathbf{v})$ for $u \in \mathbb{U}_p$. It would be slightly more convenient to work with the set $\hat{\Omega}_p = \{\mathbf{v} \in \mathbb{Q}_p^n : \|\mathbf{v}\|_p = \rho(\mathbf{v}/\|\mathbf{v}\|_p^\diamond)\}$.

We now apply Proposition 4.2 to find sequences (h_m), (h'_m) of compactly supported continuous positive functions such that $h_m \le \mathbf{1}_{\hat{\Omega}_p \times I_p} \le h'_m$ and $|h_m - h'_m|$ converges uniformly to zero as $m \to \infty$. Since h_m, h'_m are compactly supported, by definition, $\lim_{m\to\infty} L_1(h_m) = \lim_{m\to\infty} L_1(h'_m)$. Hence

$$\begin{aligned}\lim_{t\to\infty} p^{-(n-2)t}\,\mathrm{vol}(\{\mathbf{v} \in p^{-t}\hat{\Omega}_p : \mathbf{q}_p(\mathbf{v}) \in I_p\} &= \lim_{t\to\infty} p^{-(n-2)t} \int_{\mathbb{Q}_p^n} \mathbf{1}_{\hat{\Omega}_p \times I_p}(p^t\mathbf{v}, \mathbf{q}_p(v)) d\mathbf{v} \\ = \mathrm{vol}(K.\mathbf{e}_1) p^{-b} \sum_{z\in\mathbb{Z}} p^{(n-2)z} \int_K \mathbf{1}_{\hat{\Omega}_p}(p^{-z}k\mathbf{e}_1) dm(k) &=: \lambda_{\mathbf{q}_p, \hat{\Omega}_p} p^{-b}.\end{aligned}$$

Note that $\sum_{z\in\mathbb{Z}} p^{(n-2)z} \int_K \mathbf{1}_{\hat{\Omega}_p}(p^{-z}k\mathbf{e}_1) dm(k)$ is a summation over a finite set of z's. The result will now follows from the fact that $\Omega = \bigcup_{i\ge 0} p^i\hat{\Omega}$ is a disjoint union. □

Proof of Theorem 1.5. We will start by introducing two function spaces: let $\mathcal{L}$ denote the space of real-valued functions $F(\mathbf{v},\zeta)$ on $\mathbb{Q}_S^n - \{0\} \times \mathbb{Q}_S$ with support in an implicitly fixed compact set C, satisfying $F(u\mathbf{v},\zeta) = F(\mathbf{v},\zeta)$ for all $u \in \mathbb{U}_p, p \in S_f$. We equip $\mathcal{L}$ with the topology of uniform convergence on C. By $\mathcal{L}_0$, we denote the subspace of $\mathcal{L}$ consisting of the functions of the form $F(\mathbf{v},\zeta) = J_f(\|\mathbf{v}\|^\diamond,\zeta)\nu(\mathbf{v}/\|\mathbf{v}\|^\diamond)$, where $f = \prod_{p\in S} f_p$, $\nu = \prod_{p\in S}\nu_p$, f_p is continuous with a compact support on $\mathbb{Q}_{p,+}^n$ satisfying ($\mathbf{I}_f$) and ν_p is a non-negative continuous function on $\mathbb{U}_p^n$ satisfying ($\mathbf{I}_\nu$). We claim that $\mathcal{L}_0$ is dense in $\mathcal{L}$. In fact, it is easy to see from a p-adic Stone-Weierstrass Theorem (see [5]) that the approximation holds if the invariance properties are dropped from $\mathcal{L}$ and $\mathcal{L}_0$. With the invariance properties, one only needs to integrate the approximating function ν over $\prod_{p\in S}\mathbb{U}_p$ with respect to the Haar measure to obtain the desired approximating function. This implies that for each $x \in \mathcal{D}$, there are h_x, h_x' in $\mathcal{L}_0$ such that for all $\mathbf{v} \in \mathbb{Q}_S^n$ and $\zeta \in \mathbb{Q}_S$,

$$h_x(x\mathbf{v},\zeta) \le \mathbf{1}_{\hat\Omega} \le h_x'(x\mathbf{v},\zeta)$$

and

$$|L(h_x) - L(h_x')| < \epsilon.$$

where $\mathbf{1}_{\hat\Omega}$ is the characteristic function of

$$\hat\Omega = \left\{ \mathbf{v} \in \mathbb{Q}_S^n : \begin{array}{c} \rho_\infty(\mathbf{v}_\infty/\|\mathbf{v}_\infty\|_\infty)/2 < \|\mathbf{v}_\infty\|_\infty < \rho_\infty(\mathbf{v}_\infty/\|\mathbf{v}_\infty\|_\infty) \text{ and} \\ \|\mathbf{v}_p\| = \rho_p(\mathbf{v}_p/\|\mathbf{v}_p\|_p^\diamond), p \in S_f \end{array} \right\}.$$

By Proposition 5.1, Theorem 1.10, and Proposition 5.2, there exist points $x_1,\dots,x_\ell \in \mathsf{G}/\Gamma$ so that $\mathsf{H}x_i$ is closed, $1 \le i \le \ell$, and for each compact subset $\mathcal{F}$ of $\mathcal{D} - \cup_{i=1}^\ell \mathsf{H}x_i$, there exists $\mathsf{T}_0 \succ 0$ so that for all $\mathsf{T} \succ \mathsf{T}_0$ and $x = \mathsf{g}\Gamma \in \mathcal{F}$,

$$\begin{aligned}\left| \|\mathsf{T}\|^{-(n-2)} \sum_{\mathbf{v}\in\mathbb{Z}_S^n} h\left(\frac{\mathsf{g}\mathbf{v}}{\mathsf{T}^\diamond}, \mathbf{q}(\mathsf{g}\mathbf{v})\right) - L(h)\right| &\le \left| \|\mathsf{T}\|^{-(n-2)} \sum_{\mathbf{v}\in\mathbb{Z}_S^n} h\left(\frac{\mathsf{g}\mathbf{v}}{\mathsf{T}^\diamond}, \mathbf{q}(\mathsf{g}\mathbf{v})\right) - c(\mathsf{K})\int_{\mathsf{K}} \tilde f(a_\mathsf{t}\mathsf{k}x)\nu(k^{-1}\mathbf{e}_1)dm(\mathsf{k})\right| \\ &+ \left| c(\mathsf{K})\int_{\mathsf{K}} \tilde f(a_\mathsf{t}\mathsf{k}x)\nu(k^{-1}\mathbf{e}_1)dm(\mathsf{k}) - c(\mathsf{K})\int_{\mathsf{G}/\Gamma}\tilde f dg \int_{\mathsf{K}} \nu dm(\mathsf{k})\right| \\ &+ \left| c(\mathsf{K})\int_{\mathsf{G}/\Gamma}\tilde f dg\int_{\mathsf{K}}\nu dm(\mathsf{k}) - \|\mathsf{T}\|^{n-2}\int_{\mathbb{Q}_S^n} h\left(\frac{\mathsf{g}\mathbf{v}}{\mathsf{T}^\diamond}, \mathbf{q}(\mathsf{g}\mathbf{v})\right) d\mathbf{v}\right| < \epsilon\end{aligned}$$

From the definition of $L(h)$, it follows that for T large enough we have

$$\left| \|\mathsf{T}\|^{-(n-2)} \int_{\mathbb{Q}_S^n} h\left(\frac{\mathsf{g}\mathbf{v}}{\mathsf{T}^\diamond}, \mathbf{q}(\mathsf{g}\mathbf{v})\right) d\mathbf{v} - L(h)\right| \le \epsilon.$$

By approximating argument with h_x, h_x' with a suitable ϵ, for every $\theta > 0$, there exist finitely many points $x_1,\dots,x_\ell$ with $\mathsf{H}x_i$ closed and for an arbitrary compact subset $\mathcal{F} \subseteq \mathcal{D} - \cup_{i=1}^\ell \mathsf{H}x_i$, there exists $\mathsf{T}_0 \succ 0$ such that for every $x = \mathsf{g}\Gamma \in \mathcal{F}$ and every $\mathsf{T} \succ \mathsf{T}_0$, we obtain

$$(1-\theta)\mathbf{V}_{\mathsf{l},\mathbf{q},\hat\Omega}(\mathsf{T}) \le \mathbf{N}_{\mathsf{l},\mathbf{q},\hat\Omega}(\mathsf{T}) \le (1+\theta)\mathbf{V}_{\mathsf{l},\mathbf{q},\hat\Omega}(\mathsf{T}).$$

The claim will now follows from Lemma 1.2 and a standard geometric series argument. □

6. Equidistribution of translated orbits

Let G, H, and K be as before. In this section, we prove equidistribution of the translated orbit $a_{\mathsf{t}}\mathsf{K}\mathsf{g}$ in G/Γ, where $\mathsf{g} \in G$ is fixed (or varies over a compact set), and t tends to infinity. The main result will be proven in two steps. In the first step–dealt with in this section–we assume that the test function in Theorem 1.8 is bounded. The main argument, which is purely dynamical, is based on approximating the integral above, with a similar one involving orbits shifted by unipotent elements. The advantage is that the latter can be handled using an S-arithmetic version of Dani-Margulis theorem, which is generally false for unbounded functions. We will eventually need to prove Theorem 1.8 for test functions with a moderate blowup at the cusp. This step requires an analysis of certain integrals involving the α-function, which is carried out in a later section.

Theorem 6.1. *Let $n \geq 3$ be a positive integer, and let G, H, K and a_{t} be as above. Let $\phi : \mathsf{G}/\Gamma \to \mathbb{R}$ be a bounded continuous function, and q_p be a positive continuous function on $\mathbb{U}_p^n$, and $q = \prod_{p\in S} q_p$. Let $x_0 \in \mathsf{G}/\Gamma$ be such that the orbit $\mathsf{H}x_0$ is not closed. Then*

$$\lim_{\mathsf{t}\to\infty} \int_{\mathsf{K}} \phi(a_{\mathsf{t}}\mathsf{k}x_0) q(\mathsf{k}) dm(\mathsf{k}) = \int_{\mathsf{G}/\Gamma} \phi(y) d\mu(y) \int_{\mathsf{K}} q(k)\, dm(\mathsf{k}).$$

Here $d\mu$ denotes the G-invariant probability measure on G/Γ.

The proof of this theorem closely follows [3]. The integral of ϕ along the a_{t}-translates of the K-orbits is approximated by a similar integral with a_{t} replaced by the unipotent flow to which a generalized theorem of Dani and Margulis will be applied. One difference between the situation here and the one in [6] is that as $\{a_{\mathsf{t}}\}$ is a $\mathbb{Z}$-flow, the orbit needs to be "fattened up" in order to obtain rough approximations of the orbit of a unipotent flow, which is similar to intervals in $\mathbb{Q}_p$.

Fix a prime $p \in S_f$. For simplicity, let us suppress the dependence of matrices introduced below on p. For $\alpha \in \mathbb{U}_p = \{\alpha : |\alpha| = 1\}$ and $t \in \mathbb{Z}$, we set

$$d_t = \begin{pmatrix} p^{-t} & 0 \\ 0 & p^t \end{pmatrix}, \quad u_{t,\alpha} = \begin{pmatrix} 1 & \alpha p^{-t} \\ 0 & 1 \end{pmatrix}, \quad k_{t,\alpha} = \begin{pmatrix} 1 & \alpha p^t \\ -\alpha^{-1} p^t & 1 - p^{2t} \end{pmatrix},$$

$$b_{t,\alpha} = \begin{pmatrix} 1 & 0 \\ \alpha^{-1}(p^{3t} - p^t) & 1 \end{pmatrix}, \quad w_\alpha = \begin{pmatrix} 0 & -\alpha \\ \alpha^{-1} & 0 \end{pmatrix}$$

Note that $d_t = b_{t,\alpha} u_{t,\alpha} k_{t,\alpha} w_\alpha$, $\{k_{t,\alpha} : t \in \mathbb{N}, \alpha \in \mathbb{U}_p\}$ and $\{w_\alpha : \alpha \in \mathbb{U}_p\}$ embeds into K_p (with the embedding of $\mathrm{SL}_2(\mathbb{Q}_p)$ into $\mathrm{SO}(\mathbf{q})$). This is parallel to a similar formula in the real case. The only different is that w is constant in the real case, whereas here it depends on α.

Proof of Theorem 6.1. Without loss of generality, we may assume that ϕ and q are uniformly continuous. We first prove an analogous statement for $(u_{\mathsf{z}})_{\mathsf{z}\in\mathbb{Q}_S}$ instead of $(a_{\mathsf{t}})_{\mathsf{t}\in\mathsf{T}_S}$. We will then deduce the statement for the latter from a similar statement for the former by an approximation process. First, we will show that under the assumptions of Theorem 1.8, we have

Claim 1: There exist $x_1, \ldots, x_\ell \in \mathsf{G}/\Gamma$ such that the orbits $\mathsf{H}x_i$ are closed and each carries a finite H-invariant measure, such that for each compact set $\mathcal{F} \subseteq \mathcal{D} - \bigcup_{i=1}^{\ell} \mathsf{H}x_i$, for any T with $m(\mathsf{T})$ sufficiently large, and all $x \in \mathcal{F}$, we have

$$m\left\{\mathsf{k} \in \mathsf{K} : \left| \frac{1}{\lambda_S(\mathsf{I}(\mathsf{T}))} \int_{\mathsf{I}(\mathsf{T})} \phi(u_{\mathsf{z}}\mathsf{k}x)\, d\mathsf{z} - \int_{\mathsf{G}/\Gamma} \phi\, d\mu \right| > \epsilon \right\} < \epsilon.$$

Indeed, let $\mathsf{P}_1, \dots, \mathsf{P}_k$ be the subgroups provided by Theorem 1.10 for the date $(\phi, \mathsf{K}\mathcal{D}, \epsilon)$, and let $C_i \subseteq X(\mathsf{P}_i, \mathsf{U})$ be the corresponding compact sets. Define

$$Y_i = \{\mathsf{g} \in \mathsf{G} : \mathsf{Kg} \subseteq X(\mathsf{P}_i, \mathsf{U})\}.$$

Let M be the subgroup of G generated by $\mathsf{k}^{-1}\mathsf{U}\mathsf{k}$, where $\mathsf{k} \in \mathsf{K}$. The normalizer $L = N_{\mathsf{G}}(M)$ contains K, hence is of the form $L = E\mathsf{K}$ where E is a normal subgroup of H. Since $\mathsf{U} \subseteq L$ and U is not included in any proper normal subgroup of H, we obtain $M = \mathsf{H}$.

Note that for any $\mathsf{y} \in Y_i$, we have $\mathsf{k}^{-1}\mathsf{U}\mathsf{k} \subseteq \mathsf{y}\mathsf{P}_i\mathsf{y}^{-1}$, for all $\mathsf{k} \in \mathsf{K}$. So $\mathsf{H} \subseteq \mathsf{y}\mathsf{P}_i\mathsf{y}^{-1}$. It follows that there exists a subgroup of finite index P_i' in P_i such that $\mathsf{H} = \mathsf{y}\mathsf{P}_i'\mathsf{y}^{-1}$ for all $\mathsf{y} \in Y_i$. This implies that $\mathsf{y}_1^{-1}\mathsf{y}_2 \in N_{\mathsf{G}}(\mathsf{H})$ for all $\mathsf{y}_1, \mathsf{y}_2 \in Y_i$. Hence Y_i can be covered with a finite number of right cosets of H. Therefore,

$$\bigcup_{1 \le i \le k} Y_i\Gamma/\Gamma \subseteq \bigcup_{1 \le j \le l} \mathsf{H}x_j.$$

Note that $\mathsf{P}_i \cap \Gamma$ is a lattice in P_i. Since $X(\mathsf{P}_i, \mathsf{U})$ is an analytic subset of G, we have

$$m(\{\mathsf{k} \in \mathsf{K} : \mathsf{kg} \in X(\mathsf{P}_i, \mathsf{U})\gamma\}) = 0$$

for each $\mathsf{g} \in \mathsf{G} - Y_i\gamma$. Therefore

$$m\left\{\mathsf{k} \in \mathsf{K} : \mathsf{kg} \in \bigcup_{1 \le i \le k} X(\mathsf{P}_i, \mathsf{U})\Gamma\right\} = 0,$$

holds for any $\mathsf{g} \in \mathsf{G} - \bigcup_{1 \le i \le k} Y_i\Gamma$. Since $C_i \subseteq X(\mathsf{P}_i, \mathsf{U})$ and $\mathcal{F} \subseteq \mathcal{D} - \bigcup_{j=1}^{\ell} \mathsf{H}x_j$, we have

$$m\left\{\mathsf{k} \in \mathsf{K} : \mathsf{k}x \in \bigcup_{i=1}^{k} C_i\Gamma/\Gamma\right\} = 0,$$

for every $x \in \mathcal{F}$. From here it easily follows (using an analogue of Lemma 4.2. in [6]) that there exists an open subset $W \subseteq \mathsf{G}/\Gamma$ containing $\bigcup_{i=1}^{k} C_i\Gamma/\Gamma$ such that

$$m(\{\mathsf{k} \in \mathsf{K} : \mathsf{k}x \in W\}) < \epsilon$$

for any $x \in \mathcal{F}$. From the choice of $C_1, \dots, C_k$, we have that for every $y \in \mathsf{K}\mathcal{D} - W$ and all T with $m(\mathsf{T}) \gg 0$, we have

$$\left|\frac{1}{\lambda_S(\mathsf{I}(\mathsf{T}))}\int_{\mathsf{I}(\mathsf{T})} \phi(u_{\mathsf{z}}x)d\lambda_S(\mathsf{z}) - \int_{\mathsf{G}/\Gamma} \phi d\mu\right| \le \epsilon. \tag{15}$$

Let q be a bounded continuous function on K. It follows that

$$\left|\frac{1}{\lambda_S(\mathsf{I}(\mathsf{T}))}\int_{\mathsf{K}}\int_{\mathsf{I}(\mathsf{T})} \phi(u_{\mathsf{z}}x)q(\mathsf{k})d\lambda_S(\mathsf{z}) - \int_{\mathsf{G}/\Gamma} \phi d\mu \int_{\mathsf{K}} q(\mathsf{k})\, dm(\mathsf{k})\right| \le \sup_{\mathsf{k}\in\mathsf{K}} |q(\mathsf{k})|(1 + 2 \sup_{y \in \mathsf{G}/\Gamma} |\phi(y)|)\epsilon. \tag{16}$$

We will now turn to the approximation part. Fix $p \in S_f$. In what follows $d\alpha$ is the Haar measure on $\mathbb{Z}_p$ normalized such that $\int_{\mathbb{U}_p} d\alpha = 1$. First note that

$$\begin{aligned}\int_{K_p} \phi(d_t kx) q(k)\ dm(k) &= \int_{\mathbb{U}_p}\int_{K_p} \phi(b_{t,\alpha}u_{t,\alpha}k_{t,\alpha}w_\alpha kx) q(k) dm(k) d\alpha \\ &= \int_{\mathbb{U}_p}\int_{K_p} \phi(b_{t,\alpha}u_{t,\alpha}k_{t,\alpha}w_\alpha kx) q(k) dm(k_{t,\alpha}w_\alpha k) d\alpha \\ &= \int_{\mathbb{U}_p}\int_{K_p} \phi(b_{t,\alpha}u_{t,\alpha}kx) q(w_\alpha^{-1}k_{t,\alpha}k) dm(k) d\alpha.\end{aligned} \tag{17}$$

We can now write

$$\begin{aligned}&\left|\int_{K_p} \phi(d_t kx) q(k)\ dm(k) - \int_{\mathbb{U}_p}\int_{K_p} \phi(u_{t,\alpha}kx) q(w_\alpha^{-1}k)\ dm(k) d\alpha\right| \\ &= \left|\int_{\mathbb{U}_p}\int_{K_p} \phi(b_{t,\alpha}u_{t,\alpha}kx) q(w_\alpha^{-1}k_{t,\alpha}k) - \phi(u_{t,\alpha}kx) q(w_\alpha^{-1}k)\ dm(k) d\alpha\right|.\end{aligned} \tag{18}$$

Note that as $t \to \infty$, $b_{t,\alpha}$ and $k_{t,\alpha}$ converge to 1 uniformly in α. Hence, for large values of t, we have

$$\left|\int_{K_p} \phi(d_t kx) q(k)\ dm(k) - \int_{\mathbb{U}_p}\int_{K_p} \phi(u_{t,\alpha}kx) q(w_\alpha^{-1}k)\ dm(k) d\alpha\right| < \frac{\epsilon}{2}.$$

Since q is continuous, we can partition $\mathbb{U}_p$ into intervals $I_1, \dots, I_r$ such that for each $1 \le i \le r$, there exists $w_i \in K_p$ such that $|q(w_\alpha^{-1}k) - q(w_i k)| < \epsilon/4$ for every $\alpha \in I_i$. This implies that

$$\left|\int_{I_i}\int_{K_p} \phi(u_{t,\alpha}kx) q(w_\alpha^{-1}k) - \phi(u_{t,\alpha}kx) q(w_i k)\ dm(k) d\alpha\right| < \lambda_p(I_i)\epsilon/4.$$

It is clear that $I_{i,t} = \{p^{-t}\alpha : \alpha \in I_i\}$ has a measure comparable to p^t. So, for sufficiently large t, we have

$$\int_{I_i}\int_{K_p} \phi(u_{t,\alpha}kx) q(w_i k)\ dm(k) d\alpha = \int_{I_{i,t}}\int_{K_p} \phi(u_t kx) q(w_i k) d\theta(t).$$

We can now apply (15) to obtain the result.

□

7. Establishing upper bounds for the certain integrals

In this section, we will prove Theorem 1.7. In the course of the proof we will need to consider a number of integrals that are intimately connected to the integral featuring in Theorem 1.7. In the rest of this section, we fix $p \in S_f$ and denote by a_t the linear transformation a_t^p defined in equation (2): $a_t\mathbf{e}_1 = p^t\mathbf{e}_1$, $a_t\mathbf{e}_n = p^{-t}\mathbf{e}_n$, and $a_t\mathbf{e}_j = \mathbf{e}_j$, $2 \le j \le n-1$.

Lemma 7.1. *Let V be a finite dimensional vector space over $\mathbb{Q}_p$ and K be a compact subgroup of $\mathrm{GL}(V)$. Let $v \in V$ and W be a proper subspace of V such that for any finite index subgroup K' of K and any non-zero subspace $W' \subset W$, $K'W' \not\subseteq W'$. Then the subset*

$$\mathrm{tran}(v, W) = \{k \in K : kv \in W\} \subseteq K \tag{19}$$

has Haar measure zero.

Proof. Denote the Haar measure on K by m. Since K is compact, $m(K) < \infty$. Suppose that $m(\mathrm{tran}(v, W)) > 0$. Take $W' \subseteq W$ of the smallest dimension such that $\mathrm{tran}(v, W')$ has a positive measure. For each $k \in K$, set $kW' = \{kw : w \in W'\}$. For each $k' \in K$, we have $\mathrm{tran}(v, k'W') = k'\mathrm{tran}(v, W')$ and hence

$$m(\mathrm{tran}(v, k'W')) = m(\mathrm{tran}(v, W'))$$

for all $k' \in K$. If $k'W' \neq k''W'$, since $W' \cap k'^{-1}k''W'$ has dimension strictly lower than that of W', by minimality, the intersection

$$\begin{aligned}\mathrm{tran}(v, k'W') \cap \mathrm{tran}(v, k''W') =& k'\mathrm{tran}(v, W') \cap k'\mathrm{tran}(v, k'^{-1}k''W') \\ \subseteq& k'\mathrm{tran}(v, W' \cap k'^{-1}k''W')\end{aligned}$$

has measure zero.

Thus $\{kW' : k \in K\}$ is a finite set and $K' := \{k : kW' = W'\}$ is of finite index in K. $K'W' = W' \subset W'$ contradiction to the condition that $K'W' \nsubseteq W'$. □

Lemma 7.2. *Let ρ be an analytic representation of $\mathrm{SL}_n(\mathbb{Q}_p)$ on a finite dimensional normed space $(V, \| \|)$ over $\mathbb{Q}_p$ such that all elements of $\rho(\mathrm{SL}_n(\mathbb{Z}_p))$ preserve the norm on V. For $g \in \mathrm{SL}_n(\mathbb{Q}_p)$ and $v \in V$, we will denote $\rho(g)(v)$ by gv. Let K be a closed analytic subgroup of $\mathrm{SL}_n(\mathbb{Z}_p)$. Assume that V has a decomposition*

$$V = W^- \oplus W^0 \oplus W^+, \tag{20}$$

where

$$\begin{aligned} W^0 &= \{v \in V : a_t v = v, t \in \mathbb{Z}\} \\ W^\pm &= \left\{v \in V : a_t v = p^{\mp t} v, t \in \mathbb{Z}\right\}. \end{aligned}$$

Let Q be a closed subset of $\{v \in V : \|v\|_p = 1\}$. Suppose that the following conditions are satisfied:

(1) The subspace $W^- + W^0$ satisfies the assumption of Lemma 7.1;
(2) There is a positive integer ℓ such that for any nonzero $v \in W^-$,

$$\mathrm{codim}\left\{x \in \mathrm{Lie}(K) : xv \in W^-\right\} \geq \ell. \tag{21}$$

Then for any s, $0 < s < \ell$,

$$\lim_{t \to +\infty} \sup_{v \in Q} \int_K \frac{dm(k)}{\|a_t k v\|^s} = 0,$$

where m is the normalized Haar measure on K.

Proof. Let $p : V \to W^+ \oplus W^0$ and $p^+ : V \to W^+$ denote the natural projections associated to the decomposition (20). For any $v \in V$ and $r \geq 0$, define

$$\begin{aligned} D(v, r) &= \{k \in K : \|p(kv)\| \leq r\} \\ D^+(v, r) &= \left\{k \in K : \|p^+(kv)\| \leq r\right\}. \end{aligned}$$

We will show that the measure of $D(v, r)$ is bounded by Cr^ℓ, where C is uniform over all $v \in Q$. Recall that the Lie algebra $\mathrm{Lie}(K)$ is defined as the tangent space to the p-adic analytic manifold K at the identity. Consider the map $f_v : K \to W^+ \oplus W^0$ defined by $f_v(k) = p(kv)$. Since K acts by linear transformations, the derivative of f_v at the identity is given by

$$d_e f_v(x) = p(xv), \quad x \in \mathrm{Lie}(K) = T_e K \subseteq \mathfrak{sl}_n(\mathbb{Q}_p). \tag{22}$$

For every $v \in V$, set

$$L_v = \{x \in \mathrm{Lie}(K) : xv \in W^-\} = \ker d_e f_v.$$

For $k \in K$, also define the map $m_k : K \to K$ by $m_k(k') = k'k$. Note that $m_k(e) = k$ and $d_e m_k : T_e K \to T_k K$ is an isomorphism. Differentiating $f_{kv} = f_v \circ m_k$ yields $d_e f_{kv} = d_k(f_v) \circ d_e m_k$, showing that $\operatorname{rank} d_e f_{kv} = \operatorname{rank} d_k f_v$. From the assumption, if $f_v(k) = 0$, then $kv \in W^-$, hence by the assumption of the theorem, $\operatorname{codim} \ker d_e f_{kv} \geq \ell$, which implies that $\operatorname{codim} \ker d_k f_v \geq \ell$.

Let us denote by $\mathbb{P}(V)$ the projective space on V, and by $[v]$ the image of the non-zero vector v in $\mathbb{P}(V)$. We know that if $f : U \to V$ is a p-adic analytic function, then the map $x \mapsto \operatorname{rank} d_x f$ is lower semi-continuous, that is, $\liminf_{y \to x} \operatorname{rank} d_y f \geq \operatorname{rank} d_x f$. This implies that for every $v \in W^-$, there exists an open subset U containing $[v] \in \mathbb{P}(V)$ such that for $[w] \in U$, we have $\operatorname{codim} \ker d_k f_w \geq \ell$. By compactness, there exists $\epsilon > 0$ such if $\|p(kv)\| \leq \epsilon \|v\|$, then the map f_v has a rank at least ℓ. Using the implicit function theorem for local fields (e.g. [21]), for any $k \in K$ for which $\|p(kv)\| \leq \epsilon \|v\|$, we can choose local coordinate systems for $\mathcal{U}_k \subset K$ at k and for $W^+ \oplus W^0$ in which f_v is given by

$$f_v(u_1, \ldots, u_m) = (u_1, \ldots, u_\ell, \psi_{\ell+1}, \ldots, \psi_{m'}),$$

where $m = \dim K$, $m' = \dim W^0 + \dim W^+$ and $\psi_j(u_1, \ldots, u_m)$ is an analytic function for $j = \ell + 1, \ldots, m'$ depending smoothly on v. If $r \leq \epsilon$, for such a neighborhood $\mathcal{U}_k$, $k \in D(v, r)$, there is a nonnegative $\alpha = \alpha(k)$ satisfying that;

$$m\left(D(v, r) \cap \mathcal{U}_k\right) \sim \left(\frac{r}{\epsilon}\right)^{\ell + \alpha} m\left(D(v, \epsilon) \cap \mathcal{U}_k\right)$$

so that

$$\frac{m\left(D(v, r)\right)}{r^\ell} \leq \frac{1}{\epsilon^\ell} m\left(D(v, r)\right).$$

If $r \geq \epsilon$, since $m(D(v, r)) \leq 1$,

$$\frac{m\left(D(v, r)\right)}{r^\ell} \leq \frac{1}{\epsilon^\ell}.$$

By compactness of the ball $\{v \in V : \|v\| = 1\}$, we have

$$C := \sup_{\|v\| = 1, r > 0} \frac{m\left(D(v, r)\right)}{r^\ell} < \infty. \tag{23}$$

We will now have to consider the set

$$D^+(v, 0) = \{k \in K : p^+(kv) = 0\} = \{k \in K : kv \in W^0 \oplus W^-\}.$$

By Lemma 7.1, we have $m(D^+(v, 0)) = 0$. We claim that this and the compactness of Q imply that

$$\lim_{r \to 0} \sup_{v \in Q} m(D^+(v, r)) = 0. \tag{24}$$

Otherwise, pick a sequence $r_n \to 0$, and $v_n \in Q$ such that $m(D^+(v_n, r_n)) > \epsilon$ for some ϵ. Upon passing to a subsequence, we can assume that v_n converges to v.

$$\epsilon \leq m(\{k : \|p^+(kv_n)\| \leq r_n\}) \subseteq m(\{k : \|p^+(kv)\| \leq r_n + \|v_n - v\|\}).$$

Since $r_n + \|v_n - v\| \to 0$, we obtain $m(D^+(v, 0)) \geq \epsilon$, which is a contradiction.

Since $a_t|_{W^0} = Id_{W^0}$ and (23), we see that for a positive integer t, $\|a_t kv\|^s \geq \|p(kv)\|^s$ hence

$$\int_{D(v,r)-D\left(v,\frac{r}{p}\right)} \frac{dm(k)}{\|a_t kv\|^s} \leq \frac{m\left(D(v,r)\right)}{(r/p)^s} \leq \left(p^{-s}C\right) r^{\ell-s},$$

for any $v \in V$, $\|v\| = 1$ and any $r > 0$. From the fact that $m\left(D(v,0)\right) = 0$,

$$\begin{aligned} \int_{D(v,r)} \frac{dm(k)}{\|a_t kv\|^s} &= \sum_{n=0}^{\infty} \int_{D(v,p^{-n}r)\setminus D(v,p^{-n-1}r)} \frac{dm(k)}{\|a_t kv\|^s} \\ &\leq \frac{C}{p^{-s}-p^{-\ell}} r^{\ell-s}. \end{aligned} \tag{25}$$

On the other hand, since $a_t|_{W^+} = p^{-t}\, Id_{W^+}$, we get that $\|a_t kv\|^s \geq p^{ts}\|p^+(kv)\|^s$ so that for any $v \in V$, $r > 0$ and $r_1 > 0$,

$$\begin{aligned} \int_{K\setminus D(v,r)} \frac{dm(k)}{\|a_t kv\|^s} &\leq \int_{K-D^+(v,r_1)} \frac{dm(k)}{\|a_t kv\|^s} + \int_{D^=(v,r_1)\setminus D(v,r)} \frac{dm(k)}{\|a_t kv\|^s} \\ &\leq p^{-ts} r_1^{-s} + m\left(D^+(v,r_1)\right) r^{-s}. \end{aligned}$$

Therefore for a given $\epsilon_0 > 0$, since $\ell - s > 0$, take $r > 0$ and $r_1 > 0$ such that $Cr^{\ell-s}/\left(p^{-s}-p^{-\ell}\right) < \epsilon_0/3$ and $m\left(D^+(v,r_1)\right) r^{-s} < \epsilon_0/3$. Then we can take $t' > 0$ such that if $t > t'$, then $p^{-ts} r_1^{-s} < \epsilon_0/3$ so that

$$\int_K \frac{dm(k)}{\|a_t kv\|^s} < \epsilon_0.$$

□

We now write V_i for the i-th exterior power $\bigwedge^i \mathbb{Q}_p^n$, and denote by ρ_i the i-th exterior power of the natural representation $\mathrm{SL}_n(\mathbb{Q}_p)$ on $\mathbb{Q}_p^n$. Recall that for an i-tuple $J = (j_1, \ldots, j_i)$, we define $\mathbf{e}_J = \mathbf{e}_{j_1} \wedge \cdots \wedge \mathbf{e}_{j_i}$. It is easy to see that V_i decomposes into a_t-eigenspaces $V_i = W_i^- \oplus W_i^0 \oplus W_i^+$, and the eigenspaces are given by

$$\begin{aligned} W_i^- &= \mathrm{span}\left\{e_{j_1\cdots j_i} : j_1 = 1, j_i < n\right\}, \\ W_i^+ &= \mathrm{span}\left\{e_{j_1\cdots j_i} : 1 < j_1, j_i = n\right\}, \\ W_i^0 &= \mathrm{span}\left\{e_{j_1\cdots j_i} : j_1 = 1, j_i = n \text{ or } j_1 > 1, j_i < n\right\} \end{aligned}$$

Proposition 7.3. *Let $\mathbf{q}_p$ be the standard quadratic form on $\mathbb{Q}_p^n$, $n \geq 4$. Let $K = \mathrm{SL}_n(\mathbb{Z}_p) \cap \mathrm{SO}(\mathbf{q}_p)$. Let $A = A_p$ and let V_i, W_i^-, W_i^0 and W_i^+ be as above. For each V_i, define*

$$F(i) = \left\{\mathbf{x}_1 \wedge \mathbf{x}_2 \wedge \cdots \wedge \mathbf{x}_i : \mathbf{x}_1, \mathbf{x}_2, \cdots, \mathbf{x}_i \in \mathbb{Q}_p^n\right\} \subset V_i \tag{26}$$

and $Q_i = F(i) \cap \{v \in V_i : \|v\|_p = 1\}$. Then for any s, $0 < s < 2$,

$$\lim_{t\to\infty} \sup_{v\in Q_i} \int_K \frac{dm(k)}{\|a_t kv\|^s} = 0. \tag{27}$$

Proof. Let us first consider the special case of $n = 4$, in which $\mathbf{q}_p$ is given by $\mathbf{q}_p(x_1, x_2, x_3, x_4) = x_1x_4 + \alpha_2 x_2^2 + \alpha_3 x_3^2$, where $\alpha_i \in \{\pm 1, \pm u_0, \pm p, \pm u_0 p\}$. Here $u_0 \in \mathbb{U}_p$ is a representative of a quadratic non-residue in the quotient $\mathbb{Z}_p/p\mathbb{Z}_p$. Define the following subgroups of K:

$$\mathbf{U} = \left\{ u_z = \begin{pmatrix} 1 & 0 & 0 & 0 \\ z & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -\alpha_2 z^2 & -2\alpha_2 z & 0 & 1 \end{pmatrix} : z \in \mathbb{Q}_p \right\} \tag{28}$$

$$\mathbf{U}' = \left\{ u'_z = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ z & 0 & 1 & 0 \\ -\alpha_3 z^2 & 0 & -2\alpha_3 z & 1 \end{pmatrix} : z \in \mathbb{Q}_p \right\}. \tag{29}$$

Direct calculation yields

$$u_z \mathbf{e}_{123} = \mathbf{e}_{123} + 2\alpha_2 z\ \mathbf{e}_{134} + \alpha_2 z^2\ \mathbf{e}_{234}, \quad u'_z \mathbf{e}_{123} = \mathbf{e}_{123} - 2\alpha_3 z \mathbf{e}_{124} + \alpha_3 z^2 \mathbf{e}_{234},$$

Similarly, for $\mathbf{v} = x_{12}\mathbf{e}_{12} + x_{13}\mathbf{e}_{13} + x_{23}\mathbf{e}_{23}$, we have

$$u_z \mathbf{v} = x_{12}\mathbf{e}_{12} + 2\alpha_2 z x_{12}\mathbf{e}_{14} - \alpha_2 z^2 x_{12}\mathbf{e}_{24} + x_{13}\mathbf{e}_{13} + (z x_{13} + x_{23})\mathbf{e}_{23} + \alpha_2(z^2 x_{13} + 2z x_{23})\mathbf{e}_{34},$$

$$u'_z \mathbf{v} = x_{12}\mathbf{e}_{12} - (z x_{12} + x_{23})\mathbf{e}_{23} + \alpha_3(z^2 x_{12} - 2t_3 x_{23})\mathbf{e}_{24} + x_{13}\mathbf{e}_{13} - 2\alpha_3 z x_{13}\mathbf{e}_{14} - \alpha_3 z^2 x_{13}\mathbf{e}_{34}.$$

Note that if K' is a finite index subgroup of K, then it is open in K and K' always intersects $\mathbf{U}$ and $\mathbf{U}'$. The above computation shows that the subspace $W_i^- + W_i^0$ of V_i, for any i, satisfies the assumption of Lemma 7.1 with respect to K. Moreover, from the fact that subgroups $\mathbf{U}$ and $\mathbf{U}'$ are contain we obtain the condition (21) at least for $\ell = 2$ of Lemma 7.2. By Lemma 7.2, for $0 < s < 2$, we can see that

$$\lim_{t\to\infty} \sup_{v \in Q_i} \int_K \frac{dm(k)}{\|a_t k v\|^s} = 0,$$

for any i. Now let us show that (27) holds for general quadratic forms when $n \geq 4$.

For a general standard quadratic form $\mathbf{q}_p$ on $\mathbb{Q}_p^n$, for any j_2, j_3 fixed, consider the embedding $\iota(j_2, j_3)$ from $\mathrm{SO}(x_1x_4 + \alpha_{j_2}x_2^2 + \alpha_{j_3}x_3^3)$ to $\mathrm{SO}(\mathbf{q}_p)$ induced by the following injection from $\mathbb{Q}_p^4$ into $\mathbb{Q}_p^n$;

$$\mathbf{e}_1 \mapsto \mathbf{e}_1,\ \mathbf{e}_2 \mapsto \mathbf{e}_{j_2},\ \mathbf{e}_3 \mapsto \mathbf{e}_{j_3},\ \mathbf{e}_4 \mapsto \mathbf{e}_n.$$

Clearly, the embedding $\iota(j_2, j_3)$ maps subgroups $\mathbf{U}(\mathbb{Z}_p)$ and $\mathbf{U}'(\mathbb{Z}_p)$ into subgroups of the maximal compact subgroup K of $\mathrm{SO}(\mathbf{q}_p)$. For any $v \in W_i^- + W_i^0$, we can find an appropriate embedding $\iota(j_2, j_3)$ so that orbits $\iota(j_2, j_3)(\mathbf{U}(\mathbb{Z}_p)).v$ and $\iota(j_2, j_3)(\mathbf{U}'(\mathbb{Z}_p)).v$ escape the subspace $W_i^- + W_i^0$ except the identity. Therefore by the same argument with the case $n = 4$, we can come to a conclusion (27).

Note that if K' is a finite index subgroup of K, then it is open in K and K' always intersects $\mathbf{U}$ and $\mathbf{U}'$. The above computation shows that the subspace $W_i^- + W_i^0$ of V_i, for any i, satisfies the assumption of Lemma 7.1 with respect to K. Moreover, from the fact that subgroups $\mathbf{U}$ and $\mathbf{U}'$ intersect any neighbourhood of the identity, we obtain the condition (21) at least for $\ell = 2$ of Lemma 7.2. The claim will now follow from Lemma 7.2. □

In the next lemma, we will combine the results over different places $p \in S$. Recall that for $t = (t_p)_{p\in S}$, we set $a_{\mathfrak{t}} = (a_{t_p})_{p\in S}$. For $p \in S$, we denote by ϵ_p the element of $\mathbb{T}$ for which $t_p = 1$ and $t_q = 0$ for $q \neq p$. Note that $p \in S_f$, $a_{\epsilon_p} = a_1^p = \mathrm{diag}\left(p, 1, \ldots, 1, p^{-1}\right)$ as an element in $\mathrm{SO}(\mathbf{q}_p)$ (or as an element embedded in $\mathrm{SO}(\mathbf{q})$).

Lemma 7.4. *Let $n \geq 4$, $\mathbf{q}$ be a non-degenerate isotropic standard quadratic form over $\mathbb{Q}_S^n$, and K be the maximal compact subgroup of $\mathrm{SO}(\mathbf{q})$ defined above. If $\mathbf{q}$ is non-exceptional, then for any $s \in (0,2)$ and any $c > 0$, there exists a positive integer m such that for every lattice Δ in $\mathbb{Q}_S^n$, and every $\mathsf{t} \in R = \{m\epsilon_p : p \in S\}$ the following inequality holds:*

$$\int_{\mathsf{K}} \alpha_i(a_{\mathsf{t}}\mathsf{k}\Delta)^s dm(\mathsf{k}) < \frac{c}{2}\alpha_i(\Delta)^s + \omega^2 \max_{0<j<\min\{n-i,i\}} \left(\alpha_{i+j}(\Delta)\alpha_{i-j}(\Delta)\right)^{s/2}. \tag{30}$$

If $S_f \neq \emptyset$, then the same assertion holds for the case that $n = 4$, and $\mathbf{q}_\infty$ has signature $(2,2)$, with $R = \{m\epsilon_p : p \in S_f\} \cup \{\epsilon_\infty + m\epsilon_p : p \in S_f\}$.

Proof. First, let us assume that $\mathbf{q}$ is non-exceptional, and let $c > 0$ be given. We will use Proposition 7.3 and the analogous statement in [6]. For each $p \in S$, one can find an integer $m_p > 0$ such that for any $v \in F(i)$, where $F(i)$ is defined in (26) with $\|v\|_p = 1$, and any $t_p > m_p$.

$$\int_{K_p} \frac{dm(k_p)}{\|a_{t_p}^p k_p v\|_p^s} < c/2.$$

Note that

$$\int_{K_p} \frac{dm(k_p)}{\|a_0^p k_p v\|_p^s} \leq 1$$

This implies that for $\mathsf{t} = m\epsilon_p$, $p \in S$, we have

$$\int_{\mathsf{K}} \frac{dm(\mathsf{k})}{\|a_{\mathsf{t}}\mathsf{k}v\|^s} = \prod_{p\in S} \int_{K_p} \frac{dm(k)}{\|a_{t_p}^p k_p v\|_p^s} < \frac{c}{2}.$$

The argument in the case where $\mathbf{q}_\infty$ is of signature $(2,2)$ is similar. In this case, one cannot find a multiple of ϵ_∞ for which the inequality holds, but using Proposition 7.3, one can choose m large enough so that

$$\int_{K_p} \frac{dm(k)}{\|a_m^p k v\|_p^s} \cdot \int_{K_\infty} \frac{dm(k')}{\|a_1^\infty k' v\|_\infty^s} < \frac{c}{2}.$$

In either case, we deduce that for every nonzero $v \in F(i)$, and $\mathsf{t} \in R$, the following holds:

$$\int_{\mathsf{K}} \frac{dm(k)}{\|a_{\mathsf{t}}\mathsf{k}v\|^s} < \frac{c}{2}\,\frac{1}{\|v\|^s}. \tag{31}$$

For a given S-lattice Δ in $\mathbb{Q}_S^n$ and each i, there exists a Δ-rational subspace L_i of dimension i satisfying that

$$\frac{1}{d(L_i)} = \alpha_i(\Delta). \tag{32}$$

By substituting a wedge product of $\mathbb{Z}_S$-generators of $L_i \cap \Delta$ for v in (31), we have that

$$\int_{\mathsf{K}} \frac{dm(\mathsf{k})}{d_{a_{\mathsf{t}}\mathsf{k}\Delta}(a_{\mathsf{t}}\mathsf{k}L_i)^s} < \frac{c}{2}\,\frac{1}{d_\Delta(L_i)^s}. \tag{33}$$

Let $\omega > 1$ be such that for all $v \in F(i)$, $1 \leq i \leq n-1$, and $\mathsf{t} \in R$ we have

$$\omega^{-1} \leq \frac{\|a_{\mathsf{t}}v\|}{\|v\|} \leq \omega. \tag{34}$$

Let Ψ_i be the set of Δ-rational subspaces L of dimension i for which

$$d_\Delta(L) < \omega^2 d_\Delta(L_i).$$

We will prove inequality (30) by distinguishing two cases. First, assume that $\Psi_i = \{L_i\}$. Then by (34), we have

$$d_{a_\mathsf{t}\mathsf{k}\Delta}(a_\mathsf{t}\mathsf{k}L) = \|a_\mathsf{t}\mathsf{k}v\| \geq \omega^{-1}\|v\| = \omega^{-1}d(L) \geq \omega d(L_i) = \omega\|v'\| \geq \|a_\mathsf{t}\mathsf{k}v'\| = d_{a_\mathsf{t}\mathsf{k}\Delta}(a_\mathsf{t}\mathsf{k}L_i), \tag{35}$$

where v and v' are wedge products of $\mathbb{Z}_S$-generators of L and L_i respectively. By inequalities (33),(35) and the definition of α_i,

$$\int_K \alpha_i(a_\mathsf{t}\mathsf{k}\Delta)^s dm(\mathsf{k}) < \frac{c}{2}\alpha_i(\Delta)^s. \tag{36}$$

Now we assume that $\Psi_i \neq \{L_i\}$. Let M be an element of Ψ_i different from L, and let $\dim(M + L_i) = i + j$ for some $j > 0$. Using Lemma 3.8 and the inequality (34), we get that for any $\mathsf{k} \in \mathsf{K}$,

$$\begin{aligned}\alpha_i(a_\mathsf{t}\mathsf{k}\Delta) < \omega\alpha_i(\Delta) = \frac{\omega}{d(L_i)} &< \frac{\omega^2}{(d(L_i)d(M))^{1/2}} \\ &\leq \frac{\omega^2}{(d(L_i\cap M)d(L_i+M))^{1/2}} \leq \omega^2\left(\alpha_{i+j}(\Delta)\alpha_{i-j}(\Delta)\right)^{1/2},\end{aligned}$$

so that

$$\int_\mathsf{K} \alpha_i(a_\mathsf{t}\mathsf{k}\Delta)^s dm(\mathsf{k}) \leq \omega^2 \max_{0<j\leq\min\{n-i,i\}} \left(\alpha_{i+j}(\Delta)\alpha_{i-j}(\Delta)\right)^s. \tag{37}$$

The result follows from (36) and (37). □

7.1. **Coarsely superharmonic functions.** For $p \in S_f$, we will set

$$D_n^+ = \left\{\mathrm{diag}\left(p^{\lambda_1}, p^{\lambda_2}, \ldots, p^{\lambda_n}\right) \in \mathrm{GL}_n(\mathbb{Q}_p) : \lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_n, \lambda_i \in \mathbb{Z}\right\}.$$

Every element $g \in \mathrm{GL}_n(\mathbb{Q}_p)$ admits a Cartan decomposition

$$g = k_1(g)d(g)k_2(g), \quad k_1(g), k_2(g) \in \mathrm{GL}_n(\mathbb{Z}_p), d(g) \in D_n^+.$$

We will write $\lambda_i(g)$, $1 \leq i \leq n$ when $d(g) = \mathrm{diag}\left(p^{\lambda_1(g)}, p^{\lambda_2(g)}, \ldots, p^{\lambda_n(g)}\right)$.

Proposition 7.5. *Let U be a neighborhood of the identity element e in $\mathrm{SL}_n(\mathbb{Z}_p)$ given by;*

$$U = \{g \in \mathrm{SL}_n(\mathbb{Z}_p) : \|g - e\|_p < 1\}, \tag{38}$$

where $\|\cdot\|_p$ is the maximum p-norm on the vector space of $n \times n$ matrices with respect to the standard basis. Then for any $k \in U$ and $d, d' \in D_n^+$, and any $1 \leq i \leq n$, we have

$$\lambda_i(dkd') = \lambda_i(d)\lambda_i(d') \tag{39}$$

Proof. First note that $\|g\|_p = p^{-\lambda_1(g)}$. Let us denote (i,j)-entry of g by g_{ij}. Let $d = \mathrm{diag}\left(p^{\lambda_1}, \ldots, p^{\lambda_n}\right)$ and $d' = \mathrm{diag}(p^{\lambda'_1}, \ldots, p^{\lambda'_n})$ be in D_n^+. Then $dkd' = (p^{\lambda_i+\lambda'_j}k_{ij})$. Since $|k_{ii}|_p = 1$ for $1 \leq i \leq n$ and $|k_{ij}|_p < 1$ for any $i \neq j$, we obtain

$$p^{\lambda_1(dkd')} = \frac{1}{\|dkd'\|_p} = \frac{1}{|p^{\lambda_1+\lambda'_1}k_{11}|_p} = p^{\lambda_1+\lambda'_1} = p^{\lambda_1(d)}p^{\lambda_1(d')}. \tag{40}$$

Consider the representation ρ_i of $\mathrm{GL}_n(\mathbb{Q}_p)$ on the i-th exterior product $\bigwedge^i(\mathbb{Q}_p^n)$ in the usual way and denote the maximum p-norm on $\bigwedge^i \mathbb{Q}_p^n$ with respect to the standard basis by $\|\cdot\|_p$. When expressed in the standard basis of $\bigwedge^i(\mathbb{Q}_p^n)$, the entries of $\rho_i(g)$ are given by pairs (J, K), where J and K are i-element subsets of the set $\{1, 2, \ldots, n\}$. More precisely, $\rho_i(g) = (\det g_{JK})$, where g_{JK} is the $i \times i$ minor of g formed by the rows in J and columns in K. Note that the

p-norm on $\bigwedge^i \mathbb{Q}_p^n$ is also invariant under the action of $\mathrm{GL}_n(\mathbb{Z}_p)$. For $k \in U$, since $|k_{ii}|_p = 1$ and $|k_{ij}| < 1$ for $i \neq j$, we readily see that $|\det(k_{JK})| = 1$ if $J = K$ and $|\det(k_{JK})| < 1$, otherwise. This implies that

$$\|\rho_i(g)\|_p = \|\rho_i(dkd')\|_p = p^{-(\lambda_1(g)+\cdots+\lambda_i(g))}.$$

Hence we obtain

$$p^{\lambda_1(dkd')+\cdots+\lambda_i(dkd')} = \frac{1}{\|\rho_i(dkd')\|_p} = p^{\sum_{j=1}^i \lambda_j + \sum_{j=1}^i \lambda'_j} = p^{\lambda_1(d)+\cdots\lambda_i(d)} p^{\lambda_1(d')+\cdots+\lambda_i(d')}. \tag{41}$$

□

We remark that the above proposition also holds for

$$D_-^n = \left\{ \mathrm{diag}\left(p^{\lambda_1}, p^{\lambda_2}, \ldots, p^{\lambda_n}\right) \in \mathrm{GL}_n(\mathbb{Q}_p) : \lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n, \lambda_i \in \mathbb{Z} \right\},$$

instead of D_+^n.

Corollary 7.6. *Let H be a simply connected simple algebraic group in $\mathrm{GL}_n(\mathbb{Q}_p)$ and K the maximal compact subgroup $\mathrm{SL}_n(\mathbb{Z}_p) \cap H$ of H. Let $a_t = \mathrm{diag}(p^{-t}, 1, \ldots, 1, p^t)$ for $t \in \mathbb{Z}$. Then there is a neighborhood U of e in K such that for $t, s \geq 0$, we have*

$$a_t U a_s \subset K a_t a_s K.$$

Proof. The Cartan decomposition of H(see Theorem 3.14 in [16]) provides that for each $g \in H$, there are elements $k_1(g)$ and $k_2(g)$ in K such that

$$a_t g a_s = k_1(g) \mathrm{diag}\left(p^{\lambda_1(a_t g a_s)}, \ldots, p^{\lambda_n(a_t g a_s)}\right) k_2(g).$$

Take a neighborhood U of K satisfying the condition of Proposition 7.5. Then

$$\mathrm{diag}\left(p^{\lambda_1(a_t g a_s)}, \ldots, p^{\lambda_n(a_t g a_s)}\right) = \mathrm{diag}\left(p^{\lambda_1(a_t a_s)}, \ldots, p^{\lambda_n(a_t a_s)}\right) = a_t a_s,$$

so we get the result. □

Proposition 7.7. *Let $\mathsf{H} = \mathrm{SO}(\mathbf{q})$, $\mathsf{K} = \mathrm{SL}_n(\mathbb{Z}_p) \cap \mathsf{H}$ and let $\mathcal{F}$ be a family of strictly positive functions on H having the following properties:*

(a) For every $\lambda > 1$, there exists a neighborhood $V(\lambda), \lambda > 1$ of the identity in H such that for all $f \in \mathcal{F}$, $\lambda^{-1} f(\mathsf{h}) < f(\mathsf{gh}) < \lambda f(\mathsf{h})$ for any $\mathsf{h} \in \mathsf{H}$ and $\mathsf{g} \in V(\lambda)$.

(b) There exists a constant $C > 0$ such that for all $p \in S_f$, and all $\mathsf{h} \in \mathsf{H}$, we have

$$f(a_{\epsilon_p} \mathsf{h}) < C f(\mathsf{h}),$$

where a_{ϵ_p} denotes the element $\mathrm{diag}(p, 1, \ldots, 1, p^{-1})$.

(c) The functions $f \in \mathcal{F}$ are left K-invariant, that is, $f(\mathsf{kh}) = f(\mathsf{h})$, for $\mathsf{k} \in \mathsf{K}$ and $\mathsf{h} \in \mathsf{H}$.

(d) $\sup_{f \in \mathcal{F}} f(1) < \infty$.

Then there exists $0 < c = c(\mathcal{F}) < 1$ such that for any $\mathsf{t}_0 > 0$ and $b > 0$ there exists $B = B(\mathsf{t}, b) < \infty$ with the following property: If $f \in \mathcal{F}$ and

$$\int_{\mathsf{K}} f(a_{\mathsf{t}_0} \mathsf{kh}) dm(\mathsf{k}) < c f(\mathsf{h}) + b \tag{42}$$

for any $\mathsf{h} \in \mathsf{H}$, then

$$\int_{\mathsf{K}} f(a_{\mathsf{t}} \mathsf{k}) dm(k) < B$$

for all $\mathsf{t} > 0$ in the T-cone generated by R.

Proof. First note that we can replace each $f \in \mathcal{F}$ with its right K-average

$$\tilde{f}(\mathsf{h}) = \int_{\mathsf{K}} f(\mathsf{hk}) dm(\mathsf{k})$$

and $\tilde{\mathcal{F}} = \{\tilde{f} : f \in \mathcal{F}\}$ still satisfies properties (a), (b) and (c). Hence, without loss of generality, we may and will assume that for all $f \in \mathcal{F}$ and $\mathsf{h} \in \mathsf{H}$, we have

$$f(\mathsf{KhK}) = f(\mathsf{h}). \tag{43}$$

We need to prove

$$\sup_{\mathsf{t}>0} f(a_{\mathsf{t}}) < B. \tag{44}$$

By Corollary 7.6 and Lemma 5.11 in [6], we can take a neighborhood U of the identity in H such that

$$a_{\mathsf{t}_0} \mathsf{U} a_{\mathsf{t}} \in \mathsf{KV} a_{\mathsf{t}_0} a_{\mathsf{t}} \mathsf{K}$$

for any $\mathsf{t}_0, \mathsf{t} \geq 0$. From (a) and (43), it follows that

$$\int_{\mathsf{K}} f(a_{\mathsf{t}_0} \mathsf{k} a_{\mathsf{t}}) dm(\mathsf{k}) \geq \int_{\mathsf{U}\cap\mathsf{K}} f(a_{\mathsf{t}_0} \mathsf{k} a_{\mathsf{t}}) dm(\mathsf{k}) > \frac{1}{2} m_{\mathsf{K}}(\mathsf{U} \cap \mathsf{K}) f(a_{\mathsf{t}_0} a_{\mathsf{t}}). \tag{45}$$

Suppose that for all $\mathsf{t}_0 \in R$ and $b > 0$, and all $\mathsf{h} \in \mathsf{H}$ we have

$$\int_{\mathsf{K}} f(a_{\mathsf{t}_0} \mathsf{kh}) dm(\mathsf{k}) < \frac{1}{4} m_{\mathsf{K}}(\mathsf{U} \cap \mathsf{K}) f(\mathsf{h}) + b \tag{46}$$

Then by taking $\mathsf{h} = a_{\mathsf{t}}$ in (46) we see that for any $\mathsf{t} > 0$, and $\mathsf{t}_0 \in \Theta$, where Θ is the semigroup generated by R, we have

$$f(a_{\mathsf{t}_0} a_{\mathsf{t}}) < \frac{1}{2} f(a_{\mathsf{t}}) + b' \leq \max(f(a_{\mathsf{t}}), b') \tag{47}$$

where $b' = \max(2b/m_{\mathsf{K}}(U \cap \mathsf{K}), f(e))$. A simple induction shows that for any t in the semigroup generated by R, we have

$$f(a_{\mathsf{t}}) \leq b'. \tag{48}$$

Note that every t in the positive cone generated by R can be decomposed as $\mathsf{t} = \mathsf{t}_1 + \mathsf{t}_2$, where t_1 is in the semigroup generated by R and t_2 is positive and bounded in every component. Since t_2 can be written as the product of a bounded number of elements of $V(1/2)$ and ϵ_p, with $p \in S_f$, the claim follows.

□

Proof of Theorem 1.7. Define functions $f_0, f_1, \ldots, f_n$ on H by

$$f_i(\mathsf{h}) = \alpha_i(\mathsf{h}\Delta), \quad \mathsf{h} \in \mathsf{H}.$$

Since $\alpha \leq \sum_{i=0}^{n} \alpha_i$, it suffices to show that for all $0 \leq i \leq n$,

$$\sup_{\mathsf{t}>0} \int_{\mathsf{K}} f_i^s(a_{\mathsf{t}} \mathsf{k}) dm(\mathsf{k}) < \infty. \tag{49}$$

Let us check that $f_0, f_1, \ldots, f_n$ have the properties (a), (b), (c), and (d) of Proposition 7.7. First note that for $v \in F(i)$, $0 \leq i \leq n$, and $\mathsf{h} \in \mathsf{H}$, we have the trivial bound $\|\mathsf{h}v\| \leq \| \wedge^i \mathsf{h}\| \|v\|$. This proves (a) and (b). Since the action of K preserves the norm on $\bigwedge^i(\mathbb{Q}_S^n)$, each f_i is left K

invariant. It is clear that $f_i(1)$ are uniformly bounded as Δ runs over a compact set of lattices. From Lemma 7.4 with $h\Delta$, for any i, $0 < i < n$ and $h \in \mathsf{H}$, we see that

$$\int_{\mathsf{K}} f_i^s(a_{\mathsf{t}}\mathsf{k}\mathsf{h})dm(\mathsf{k}) < \frac{c}{2}f_i^2 + \omega^2 \max_{0<j<\min\{n-i,i\}} (f_{i+j}f_{i-j})^{s/2}. \tag{50}$$

Define

$$f_{\epsilon,s} = \sum_{0\le i\le n} \epsilon^{i(n-i)} f_i^s,$$

since $\epsilon^{i(n-i)} f_i^s < f_{\epsilon,s}$, $f_0 = 1$ and $f_n = d(\lambda)^{-1}$, by putting $\epsilon = c/(2n\omega^2)$, we have the inequality (42) of Proposition 7.7.

$$\begin{aligned}\int_{\mathsf{K}} f_{\epsilon,s}(a_{\mathsf{t}}\mathsf{k}\mathsf{h})dm(k) &< 1 + d(\Delta)^{-s} + \frac{c}{2}f_{\epsilon,s} + n\epsilon\omega^2 f_{\epsilon,s} \\ &= cf_{\epsilon,s} + 1 + d(\Delta)^{-1}. \end{aligned} \tag{51}$$

Let $\mathcal{C}$ be an arbitrary compact set of unimodular lattices Δ and let $\mathcal{F}$ be the family of $f_{\epsilon,s}$ as Δ runs over $\mathcal{C}$. Then $\mathcal{F}$ satisfies the conditions of Proposition 7.7. From (51) it follows that there the constants b and c can be chosen uniformly for the family $f_{\epsilon,s} \in \mathcal{F}$ constant c and b. Since $\alpha_i(h\Delta)^s \le \epsilon^{-i(n-i)} f_{\epsilon,s}(h)$, by Proposition 7.7, we conclude that there exists a constant $B > 0$ such that for each i, all $\mathsf{t} \succ 0$ and all $\Delta \in \mathcal{C}$,

$$\int_{\mathsf{K}} \alpha_i(a_{\mathsf{t}}\mathsf{k}\Delta)^s dm(\mathsf{k}) < B.$$

□

We will now use this bound to prove the main result.

$$A(r) = \{x \in G/\Gamma : \alpha(x) \le r\}. \tag{52}$$

Using Mahler's compactness criterion([11]), we see that $A(r)$ is compact for any $r > 0$.

proof of Theorem 1.8. We may assume that ϕ is nonnegative. For each $r \in \mathbb{R}_{>0}$, we choose a continuous function g_r on G/Γ such that $0 \le g_r(x) \le 1$, $g_r(x) = 1$ if $x \in A(r)$ and $g_r(x) = 0$ outside $A(r+1)$. Then

$$\phi(a_{\mathsf{t}}\mathsf{k}x) = (g_r\phi)(a_{\mathsf{t}}\mathsf{k}x) + ((1-g_r)\phi)(a_{\mathsf{t}}\mathsf{k}x).$$

Following the proof of Theorem 3.5 in [6], let $\beta = 2 - s$. Since $((1-g_r)\phi)(y) = 0$ if $y \in A(r)$,

$$\begin{aligned}((1-g_r)\phi)(y) &\le C(1-g_r)(y)\alpha(y)^{2-\beta} \le C\alpha(y)^{2-\beta/2}(1-g_r)(y)\alpha(y)^{-\beta/2} \\ &\le Cr^{-\beta/2}\alpha(y)^{2-\beta/2}.\end{aligned}$$

By Theorem 1.7 and the fact that $\|\nu\|_\infty < \infty$, there exists $C' > 0$ such that

$$\begin{aligned}\int_{\mathsf{K}} ((1-g_r)\phi)(a_{\mathsf{t}}\mathsf{k}x)\nu(\mathsf{k})dm(\mathsf{k}) &\le Cr^{-\beta/2}\int_{\mathsf{K}} \alpha(a_{\mathsf{t}}\mathsf{k}x)^{2-\beta/2}\nu(\mathsf{k})dm(\mathsf{k}) \\ &\le C'r^{-\beta/2}\end{aligned} \tag{53}$$

On the other hand, since $g_r\phi$ is compactly supported, we see that for sufficiently large $\mathsf{t} \in \mathbb{R}_{>0} \times \mathbb{N}^s$ in the cone generated by R

$$\left|\int_{\mathsf{K}} (g_r\phi)(a_{\mathsf{t}}\mathsf{k}x)\nu(\mathsf{k})dm(\mathsf{k}) - \int_{G/\Gamma}(g_r\phi)(y)dg(y)\int_{\mathsf{K}}\nu(\mathsf{k})dm(\mathsf{k})\right| < \epsilon/2, \tag{54}$$

where dg is the normalized Haar measure on G/Γ. Since $g_r\phi \to \phi$ as $r \to \infty$, (53) and (54) show the theorem. □

8. Counterexamples for ternary forms

In this section, we will prove Theorem 1.11 which partially shows that Theorem 1.5 is optimal in the sense that for some exceptional forms the asymptotics does not hold.

We will first introduce the a family of quadratic forms which will be useful for the construction. Fix $\alpha_p \in \mathbb{Q}_p$, and define

$$\mathbf{q}^{\alpha_p}(\mathbf{x}) = x_1^2 + x_2^2 - \alpha_p^2 x_3^2.$$

Similarly, for an S-vector $\alpha = (\alpha_p)_{p\in S} \in \mathbb{Q}_S$, we set $\mathbf{q}^\alpha = (\mathbf{q}^{\alpha_p})_{p\in S}$. In the case $\alpha \in \mathbb{Q}$, with a slight abuse of notation, we will also use $\mathbf{q}^\alpha$ as a shorthand for the quadratic form on $\mathbb{Q}_S^n$ associated to the constant S-vector with $\alpha_p = \alpha$ for all $p \in S$. We will first prove an S-adic version of a well-known fact from number theory adapted to fulfill our purpose.

Lemma 8.1. *Given $\alpha \in \mathbb{Q}$, there exists a constant $c_\alpha > 0$ such that for a sufficiently large S-time $\mathsf{T} = (T_p)_{p\in S}$, there exists at least $c_\alpha \|\mathsf{T}\| \log \|\mathsf{T}\|$ vectors $\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{Z}_S^3$ which satisfy*

(1) $\mathbf{q}^\alpha(\mathbf{x}) = 0$.
(2) $\|x_i\|_p = T_p$ for every $1 \le i \le 3$ and $p \in S_f$.
(3) $\|\mathbf{x}\|_\infty \le T_\infty$.

Before we proceed to the proof, we emphasize that condition (2) is a stronger form of $\|x\| = T_p$ which is needed for the following arguments.

Proof. It will be slightly more convenient to first work with the quadratic form $\mathbf{q}'(\mathbf{x}) = x_1x_2 - x_3^2$, which is $\mathbb{Q}$-equivalent to $\mathbf{q}^\alpha$ for every α. For $p \in S_f$, write $T_p = p^{n_p}$, and set $a = \prod_{p\in S_f} p^{-n_p} = T_\infty \|\mathsf{T}\|^{-1}$. We will consider triples

$$x_1 = aku^2, \quad x_2 = akv^2, \quad x_3 = akuv$$

where $u, k, v \in \mathbb{Z}$ with $\gcd(u, v) = 1$ and $\gcd(u, p) = \gcd(v, p) = \gcd(k, p) = 1$ for all $p \in S_f$. It is clear that $\mathbf{x}$ satisfies conditions (1) and (2). Moreover, the condition (3) will also be satisfied if k is chosen such that

$$|u|, |v| \le \sqrt{T_\infty/3ak}.$$

It is well-known that the density of the pairs (u, v) with $\gcd(u, v) = 1$ is $c = 6/\pi^2 > 0$. We can thus produce at least as many as

$$\sum_{\substack{1\le k\le T_\infty/3a \\ \gcd(k,p)=1, p\in S}} c\frac{T_\infty}{3ak} = c\ \|\mathsf{T}\| \sum_{\substack{1\le k\le T_\infty/3a \\ \gcd(k,p)=1,\ p\in S}} \frac{1}{k} \gg \|\mathsf{T}\| \log \|\mathsf{T}\|$$

solutions. Note that the forms $\mathbf{q}^\alpha$ can be obtained from $\mathbf{q}'$ applying a rational linear transformation A, which changes the real and p-adic norms by a bounded factor. This completes the proof. □

We will need the following lemma, which is an S-arithmetic version of Lemma 3.15 in [6]. We will assume that $\Omega = \prod_{p\in S} \mathbb{U}_p^n$ is the product of unit spheres, and drop it from the notation for the counting function.

Lemma 8.2. *Let $\mathsf{I} = \prod_{p\in S} I_p$ be an S-interval. Given $\epsilon > 0$ and an S-time T_0, the set of vectors $\beta = (\beta_p)_{p\in S} \in \mathbb{Q}_S$ for which there exists $\mathsf{T} \succ \mathsf{T}_0$ such that*

$$\mathbf{N}_{\mathsf{I},\mathbf{q}^\beta}(\mathsf{T}) \geq \|\mathsf{T}\|(\log\|\mathsf{T}\|)^{1-\epsilon}$$

is dense in $\mathbb{Q}_S$.

Proof. For the notational simplicity, we will set $I_\infty = [1/4, 1/2]$ and $I_p = \mathbb{Z}_p$ for $p \in S_f$. It will be clear that the proof works for any other choice of intervals. For $\alpha \in \mathbb{Q}$ and T large, define

$$L(\alpha, \mathsf{T}) = \{\mathbf{x} \in \mathbb{Z}_S^3 : \mathbf{q}^\alpha(\mathbf{x}) = 0,\ \|\mathbf{x}\|_\infty \in [T_\infty/2, T_\infty], \quad \|\mathbf{x}\|_p = T_p,\ \forall p \in S_f\}.$$

By Lemma 8.1, for large enough T, we have

$$\operatorname{card} L(\alpha, \mathsf{T}) \geq \frac{c_\alpha}{4}\|\mathsf{T}\|\log\|\mathsf{T}\| \geq \|\mathsf{T}\|(\log\|\mathsf{T}\|)^{1-\epsilon}.$$

Note that if $\mathbf{x} = (x_1, x_2, x_3) \in L(\alpha, \mathsf{T})$ then

$$\frac{T_\infty^2}{4(1+\alpha^2)} \leq x_3^2 \leq \frac{T_\infty^2}{1+\alpha^2}.$$

By condition (2) in Lemma 8.1, we also have $|x_3|_p = T_p = p^{m_p}$ for every $p \in S$. Set

$$\beta_\infty^2 = \alpha^2 - (1+\alpha^2)T_\infty^{-2}, \quad \beta_p^2 = \alpha^2 + u_p p^{m_p}$$

where $|u_p|_p = 1$. Note that for sufficiently large T_p, $\alpha^2 + u_p p^{m_p}$ is a square in $\mathbb{Q}_p$ and β is well-defined. A simple computation shows that for any $\mathbf{x} \in L(\alpha, \mathsf{T})$ we have

$$\begin{aligned} \mathbf{q}^{\beta_\infty}(\mathbf{x}) &= \mathbf{q}^\alpha(\mathbf{x}) + (1+\alpha^2)T_\infty^{-2}x_3^2 \in [1/4, 1], \\ \mathbf{q}^{\beta_p}(\mathbf{x}) &= \mathbf{q}^\alpha(\mathbf{x}) + u^2 x_3^2 \in \mathbb{Z}_p. \end{aligned} \tag{55}$$

This implies that the

$$\mathbf{N}_{I,\mathbf{q}^\beta}(\mathsf{T}) \geq \operatorname{card} L(\alpha, \mathsf{T}) \gg \|\mathsf{T}\|\log\|T\|^{1-\epsilon}.$$

It is easy to see that the set of β obtained in this way is dense in $\mathbb{Q}_S$.

□

We can now ready to give a proof of the main theorem. Assume that for a given T_0, we denote by $W(\mathsf{T}_0)$ the set of $\gamma \in \mathbb{Q}_S$ such that there exists $\beta \in \mathbb{Q}_S^n$ satisfying $|\beta - \gamma| < \|\mathsf{T}\|^{-3}$ and

$$\mathbf{N}_{I,\mathbf{q}^\beta}(\mathsf{T}) \geq \|\mathsf{T}\|\log\|\mathsf{T}\|^{1-\epsilon}.$$

From the definition and Lemma 8.2, it is clear that $W(\mathsf{T}_0) \subseteq \mathbb{Q}_S^n$ is an open and dense. Let T_j be a sequence of S-times going to infinity. Then $W = \cap_{j=1}^\infty W(\mathsf{T}_j)$ is a set of second category, and hence contains an irrational form. For any $\gamma \in W$, there exists an infinite sequence β^i and T_i satisfying $|\gamma_p - \beta_p^i|_p < \|\mathsf{T}_i\|^{-3}$ for all $p \in S$ and

$$\mathbf{N}_{I,\mathbf{q}^{\beta^i}}(\mathsf{T}_i) \geq \|\mathsf{T}_i\|\log\|\mathsf{T}_i\|^{1-\epsilon}.$$

Set $R_j = \|\mathsf{T}_i\|$. If $\|\mathbf{x}\|_\infty < R_j$, it is easy to see that

$$\|\mathbf{q}_p^{\beta^i}(\mathbf{x}) - \mathbf{q}_p^\gamma(\mathbf{x})\|_\infty = O(R_i^{-1}) < 1/8$$

for $i \gg 1$ and $p \in S$. This implies that $\mathbf{q}_\infty^\gamma(\mathbf{x}) \in [1/8, 2]$ and $\mathbf{q}_p^\gamma(\mathbf{x}) \in \mathbb{Z}_p$ for $p \in S_f$. The claim follows from here.

The argument when all $\mathbf{q}_p$ are equivalent to $x_1x_2 - x_3x_4$ is similar. An argument along the same lines as the one given above, using the parametrization of the solutions to $x_1x_2 - x_3x_4$ given by

$$x_1 = auv, \quad x_2 = azw, \quad x_3 = auz, \quad x_4 = avw$$

establishes the result in this case.

Appendix A. A version of Witt theorem

Proof. For simplicity, let us denote Q_p, K_p and $\mathbf{v}_p$ by Q, K and $\mathbf{v}$, etc. We need to prove that for given two vectors $\mathbf{v}_1$ and $\mathbf{v}_2$ in $\mathbb{Q}_p^n$ with the same p-norm and the same quadratic value, there is an element k in K such that $k.\mathbf{v}_1 = \mathbf{v}_2$. For convenience, we may assume that $\mathbf{v}_1 = \mathbf{e}_1$ and let $\mathbf{v}_2 = \mathbf{v} \in \mathbb{Z}_p^n$. Since the same argument will be applied to both isotropic or nonisotropic vectors, we will think of a quadratic form $Q(x_1, \dots, x_n)$ as one of

$$u_1x_1^2 + \cdots + u_ix_i^2 + p(u_{i+1}x_{i+1}^2 + \cdots + u_nx_n^2) \quad \text{or}$$
$$x_1x_2 + u_3x_3^2 + \cdots + u_ix_i^2 + p(u_{i+1}x_{i+1}^2 + \cdots + u_nx_n^2).$$

depending on the case we want to treat. Here u_i's are units in $\mathbb{Z}_p$.

Then we can write the corresponding symmetric matrix $B = B_Q$ as $\begin{pmatrix} B' & \\ & pB'' \end{pmatrix}$, where B' and B'' are nondegenerate mod p. The proposition demands to find a matrix $k \in K$ satisfying that

(a) $k.\mathbf{e}_1 = \mathbf{v}$ and
(b) ${}^tkBk = B$.

Since $\mathbb{Z}_p$ is the inverse limit of $\mathbb{Z}/p^j\mathbb{Z}$, $j \to \infty$, we will construct a chain $k^j \in \mathrm{GL}_n(\mathbb{Z}/p^{j+1}\mathbb{Z})$ such that

(a') $k^j.\mathbf{e}_1 = \mathbf{v} \bmod p^{j+1}$,
(b') ${}^tk^jBk^j = B \bmod p^{j+1}$ and
(c') $k^j = k^{j+1} \bmod p^{j+1}$.

Then the inverse limit of $(k^j)_{j=0}^{\infty}$ will be an element satisfying the conditions (a) and (b). Let us denote $k^j = k_0 + pk_1 + p^2k_2 + \cdots + p^jk_j$.

Step 1. $j = 0$. Let $k^0 = k_0 = \begin{pmatrix} X_0 & Y_0 \\ Z_0 & W_0 \end{pmatrix}$ depending on the size of B' and B''. By the condition (a'), the first column $\mathbf{v}_0$ of k_0 is given by $\mathbf{v} \bmod p$. We want to find a solution of the following equation;

$$\begin{aligned} \begin{pmatrix} B' & \\ & 0 \end{pmatrix} &= \begin{pmatrix} {}^tX_0 & {}^tZ_0 \\ {}^tY_0 & {}^tW_0 \end{pmatrix} \begin{pmatrix} B' & \\ & 0 \end{pmatrix} \begin{pmatrix} X_0 & Y_0 \\ Z_0 & W_0 \end{pmatrix} \bmod p \\ &= \begin{pmatrix} {}^tX_0B'X_0 & {}^tX_0B'Y_0 \\ {}^tY_0B'X_0 & {}^tY_0B'Y_0 \end{pmatrix} \bmod p. \end{aligned}$$

By the assumption of our quadratic form, $Q(pr_i(\mathbf{e}_1)) = Q(pr_i(\mathbf{v}_0))$, where $pr_i : (x_1, \dots, x_n) \in (\mathbb{Z}/p\mathbb{Z})^n \to (x_1, \dots, x_i) \in (\mathbb{Z}/p\mathbb{Z})^i$. Applying Theorem **??** to $\big((\mathbb{Z}/p\mathbb{Z})^i, B'\big)$, we can get an isometry X_0 satisfying that the first column is $pr_i(\mathbf{v}_0)$ and ${}^tX_0B'X_0 = B' \bmod p$. Since ${}^tX_0B'$ is invertible, we should take Y_0 as 0.Note that In this step, we can not determine Z_0 and W_0.

Step 2. $j = 1$. The matrix $k^1 = \begin{pmatrix} X_0 + pX_1 & Y_0 + pY_1 \\ Z_0 + pZ_1 & W_0 + pW_1 \end{pmatrix}$ has the first column as $\mathbf{v}_0 + p\mathbf{v}_1$ and should satisfy the following equations;

$$\begin{aligned} {}^tX_0B'X_0 + p({}^tX_0B'X_1 + {}^tX_1B'X_0 + {}^tZ_0B''Z_0) &= B' \bmod p^2 \\ {}^tX_0B'Y_0 + p({}^tX_1B'Y_0 + {}^tX_0B'Y_1 + {}^tZ_0B''W_0) &= 0 \bmod p^2 \\ {}^tY_0B'Y_0 + p({}^tY_0B'Y_1 + {}^tY_1B'Y_0 + {}^tW_0B''W_0) &= pB'' \bmod p^2 \end{aligned}$$

Since $Y_0 = 0$ and ${}^tX_0B'X_0 = B' + pC_1^{11} \bmod p^2$ for some symmetric matrix C_1^{11}, we can reduce the above equations as;

$$\begin{aligned} C_1^{11} + {}^tX_0B'X_1 + {}^tX_1B'X_0 + {}^tZ_0B''Z_0 &= 0 \bmod p \\ {}^tX_0B'Y_1 + {}^tZ_0B''W_0 &= 0 \bmod p \\ {}^tW_0B''W_0 &= B'' \bmod p \end{aligned}$$

Take any Z_0 such that the first column is $(v_{i+1}, \ldots, v_n)$ and any W_0 satisfying ${}^tW_0B''W_0 = B'' \bmod p$ using Witt theorem. Then it suffices to show that there is a matrix X_1 with the given first column satisfying the equation $C_1 + {}^tX_0B'X_1 + {}^tX_1B'X_0 + {}^tZ_0B''Z_0 = 0 \bmod p$.

Step 3. We claim that for a given invertible symmetric matrix A and any symmetric matrix C, there is a solution X of the equation ${}^tXA + AX = C$.

By considering the space of n by n matrices as a n^2- dimensional vector space, we can rewrite the above equation by;

$$\begin{pmatrix} A_{11} & A_{12} & \cdots & A_{1n} \\ A_{21} & A_{22} & \cdots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{n1} & A_{n2} & \cdots & A_{nn} \end{pmatrix} \begin{pmatrix} [X]^1 \\ [X]^2 \\ \vdots \\ [X]^n \end{pmatrix} = \begin{pmatrix} [C]^1 \\ [C]^2 \\ \vdots \\ [C]^n \end{pmatrix}, \tag{56}$$

where a block matrix A_{ij} is defined by

$$A_{ij} = \begin{cases} \begin{pmatrix} a_{11} & a_{21} & \cdots & \cdots & a_{n1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 2a_{1i} & 2a_{2i} & \cdots & \cdots & 2a_{ni} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{1n} & a_{2n} & \cdots & \cdots & a_{nn} \end{pmatrix} & \text{when } A = (a_{st}), \quad i = j; \\ \text{All entries are zero except } j\text{-th row is} \\ \qquad\qquad (a_{1i}, a_{2i}, \ldots, a_{ni}), \quad i \neq j. \end{cases}$$

Since ${}^tXA + AX$ and C are both symmetric, after removing rows repeated(for example, one of second row and (n+1)th row corresponding to c_{12} and c_{21}) we get a linear equation from $(\mathbb{Z}/p\mathbb{Z})^{n^2}$ to $(\mathbb{Z}/p\mathbb{Z})^{n(n+1)/2}$. Furthermore, from the fact that A is invertible, the rank of this reduced linear equation is exactly $n(n+1)/2$ which tells us that there is a solution X.

Now let us show that $C_1 + {}^tX_0B'X_1 + {}^tX_1B'X_0 + {}^tZ_0B''Z_0 = 0 \bmod p$ has a solution when C_1, B', X_0, B'', Z_0 and the first column of X_1 are given. That is, the first n^2 by n submatrix ${}^t(A_{11}, \ldots, A_{n1})$ of the equation (1) is erased together with the n variables $[X]^1$.

Before that, let us assume that the removed row among the repeated rows in the above argument is always the former one. That is, in the example, we will remove the second row and leave $(n+1)$th row. Hence the entries of the $n(n+1)/2$ by n submatrix of the reduced matrix are zero except the first row, and consequently the rank of the linear equation $C_1 +^t X_0B'X_1 +^t X_1B'X_0 +^t Z_0B''Z_0 = 0 \bmod p$ is $n(n+1)/2-1$. On the other hand c_{11} in the equation (56) can be also removed since it is determined by A_{11} and $[X]^1$. Therefore if we check that the (1,1)-entry of $C_1 +^t X_0B'X_1 +^t X_1B'X_0 +^t Z_0B''Z_0 = 0 \bmod p$ holds, we can find a required matrix X_1. However this follows from the fact that $Q(\mathbf{e}_1) = Q(\mathbf{v}) = Q(\mathbf{v}_0+p\mathbf{v}_1) \bmod p$.

Step 4. In general, suppose that there exists a solution $k = k_0 + pk_1 + p^2k_2 + \cdots + p^nk_n + \cdots$ satisfying (a) and (b). Then from the condition ${}^tkBk = B$,

$$\begin{aligned}\begin{pmatrix} B' & \\ & pB'' \end{pmatrix} &= \sum_{j=0}^{\infty} p^j \left(\sum_{i=0}^{j} \begin{pmatrix} {}^tX_i & {}^tZ_i \\ {}^tY_i & {}^tW_i \end{pmatrix} \begin{pmatrix} B' & \\ & pB'' \end{pmatrix} \begin{pmatrix} X_{j-i} & Y_{j-i} \\ Z_{j-i} & W_{j-i} \end{pmatrix} \right) \\ &= \sum_{j=0}^{\infty} p^j \left(\sum_{i=0}^{j} \begin{pmatrix} {}^tX_iB'X_{j-i} & {}^tX_iB'Y_{j-i} \\ {}^tY_iB'X_{j-i} & {}^tY_iB'Y_{j-i} \end{pmatrix} \right) + \\ &\qquad \sum_{j=0}^{\infty} p^{j+1} \left(\sum_{i=0}^{j} \begin{pmatrix} {}^tZ_iB''Z_{j-i} & {}^tZ_iB''W_{j-i} \\ {}^tW_iB''Z_{j-i} & {}^tW_iB''W_{j-i} \end{pmatrix} \right).\end{aligned}$$

Hence we should find X_j, Y_j, Z_{j-1} and W_{j-1} inductively. Take any Z_{j-1} with the first column ${}^t(v_{j-1}^{i+1}, \ldots, v_{j-1}^n)$, where $\mathbf{v} = \mathbf{v}_0 + p\mathbf{v}_1 + \cdots + p^j\mathbf{v}_j + \cdots$. Then by step3 with the fact that $B(\mathbf{v},\mathbf{v}) = \sum_{k=0}^{j} p^j \left(\sum_{i=0}^{k} B(\mathbf{v}_i, \mathbf{v}_{k-i}) \right) \bmod p^{j+1}$, we can find an appropriate X_j, W_{j-1} and Y_j satisfying the following equations.

$$\begin{aligned} {}^tX_0B'X_j +^t X_jB'X_0 &= -\sum_{i=1}^{j-1} {}^tX_iB'X_{j-i} - \sum_{i=0}^{j-1} {}^tZ_iB''Z_{j-i-1} + C_j^{11} \mod p, \\ {}^tW_{j-1}B''W_0 +^t W_0B''W_{j-1} &= -\sum_{i=1}^{j-1} \left({}^tW_iB''W_{j-i-1} +^t Y_iB'Y_{j-i} \right) + C_n^{22} \mod p, \\ {}^tX_0B'Y_j &= -\sum_{i=0}^{j-1} \left({}^tX_iB'Y_{j-i} +^t Z_iB''W_{j-i-1} \right) + C_j^{12} \mod p, \end{aligned}$$

where C_j^{11}, C_j^{22} and C_j^{12} are obtained from the equations of the formal level $j-1$(See the step2).

Consequently, we can find $k \in GL(n, \mathbb{Z}_p)$ such that ${}^tkBk = B$. Since $\det k = \pm 1$, k may not be an element of K. However we can easily find $k' \in K$ from k with $k'.\mathbf{e}_1 = \mathbf{v}$ using reflections in $\mathbb{Q}_p^n$. □

References

[1] Borel, Prasad, *Values of isotropic quadratic forms at S-integral points.* Compositio Math. 83(1992), no.3, 347-372.

[2] H. Davenport, *Analytic methods for Diophantine equations and Diophantine inequalities.* Second edition. With a foreword by R. C. Vaughan, D. R. Heath-Brown and D. E. Freeman. Edited and prepared for publication by T. D. Browning. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2005. xx+140 pp.

[3] S. Dani and G. Margulis, *Limit distributions of orbits of unipotent flows and values of quadratic forms*, I. M. Gel'fand Seminar, 91-137, Adv. Soviet Math., 16, Part 1, Amer. Math. Soc., Providence, RI, 1993.
[4] H. Davenport and D. Ridout, *Indefinite quadratic forms*, Proc. London Math. Soc. (3) 9 1559, 544-555.
[5] J. Dieudonne, Sur les functions continues p-adiques, Bull. Sci. Math. (2) 68 (1994), 79-95.
[6] A. Eskin, G. Margulis and S. Mozes, *Upper bounds and asymptotics in a quantitative version of the Oppenheim conjecture*, Ann. of Math. (2) 147 (1998), no. 1, 93-141.
[7] A. Eskin, G. Margulis and S. Mozes, *Quadratic forms of signature* $(2, 2)$ *and eignevalue spacings on rectangular* 2*-tori.* Ann. of Math. (2) 161(2005), no. 2, 679-725.
[8] G. H. Hardy, Ramanujan : Twelve Lectures on subsects suggested by his life and work, Chelsea Publishing Company, New York 1959 iii+236 pp.
[9] G.H. Hardy, On the Expression of a Number as the Sum of Two Squares, Quart. J. Math. 46, (1915), pp.263?283.
[10] M.N. Huxley, *Integer points, exponential sums and the Riemann zeta function,* Number theory for the millennium, II (Urbana, IL, 2000) pp.275?290,
[11] D. Kleinbock and G. Tomanov, *Flows on S-arithmetic homogeneous spaces and applications to metric Diophantine approximation*, Comment. Math. Helv. 82(2007), no. 3, 519-581.
[12] Landau, E. *Neue Untersuchungen uber die Pfeiffer'sche Methode zur Abschatzung von Gitterpunktanzahlen.* Sitzungsber. d. math-naturw. Classe der Kaiserl. Akad. d. Wissenschaften, 2. Abteilung, Wien, No. 124, 469-505, 1915.
[13] K. Mallahi Karai, Distribution of orbits of unipotent groups on S-arithmetic homogeneous spaces, preprint. arXiv:submit/1554931
[14] G. A. Margulis, *Formes quadratriques indéfinies et flots unipotents sur les espaces homogénes.* (French summary), C. R. Acad. Sci. Paris. Sér. I Math. 304 (1987), no. 10. 249-253.
[15] A. Oppenheim, *The minima of indefinite quaternary quadratic forms*, Proc. Nat. Acad. Sci. U. S. A. 15 (9), 724-727.
[16] V. Platonov, A. Rapinchuk, *Algebraic Groups and Number Theory* , Academic Press, INC., 1994.
[17] M. Ratner, *Raghunathan's conjectures for Catesian products of real and p-adic Lie groups*, Duke Math. J. 77 (1995), no. 2, 275-382.
[18] D. Ridout, *Indefinite quadratic forms*, Mathematika, 5, 1958, 122-124.
[19] W. Schmidt, *Approximation to algebraic numbers,* Enseignement Math. (2) 17 (1971), 187-253.
[20] J.-P. Serre, *A course in Arithmetic.* Title of the French original edition : Cours d'Arithmétique. Graduate Texts in Mathematics Vol. 7 (New York, Springer-Verlag, 1973).
[21] J.-P. Serre, *Lie Algebras and Lie Groups.* Lectures Notes in Mathematics, Vol. 1500. (Springer-Verlag Berlin Heidelberg New York, 1965).
[22] J.-P. Serre, *Quelques applications du théoréme de densité de Chebotarev*, Inst. Hautes. Études Sci. Publ. Math., No. 54 (1981), 323-401.
[23] G. Tomanov, *Orbits on Homogeneous Spaces of Arithmetic Origin and Approximations*, Adv. Stud. Pure Math., 26, Math. Soc. Japan, Tokyo, 2000.